

Acridotheres tristis  [简体中文](#) [正體中文](#)

System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Aves	Passeriformes	Sturnidae

Common name house myna (English), common myna (English), Calcutta myna (English), mynah (English), Hirtenmaina (German), German Indischer mynah (English), manu rataro (English, Cook Islands), manu kavamani (English, Cook Islands), manu teve (English, Cook Islands), piru (English, Cook Islands), manu kaomani (English, Cook Islands), manu (English, Cook Islands), talking myna (English), Martin triste (French), Indian mynah (English), Indian myna (English)

Synonym *Acridotheres tristis* , (Linnaeus, 1766)

Similar species *Manorina melanocephala*, *Manorina flavigula*

Summary The common myna (*Acridotheres tristis*), also called the Indian myna, is a highly commensal Passerine that lives in close association with humans. It competes with small mammals and bird for nesting hollows and on some islands, such as Hawaii and Fiji, it preys on other birds' eggs and chicks. It presents a threat to indigenous biota, particularly parrots and other birdlife, in Australia and elsewhere.



[view this species on IUCN Red List](#)

Species Description

Indian mynas are 23 to 26 cm long, weigh 82 to 143 g and have a wing-span of 120 to 142 mm (Markula Hannan-Jones & Csurhes 2009). The common myna has a medium to heavy build and a cocoa brown colour (Massam 2001). The head, neck and upper breast of the adult is glossy black, while the undertail coverts, tail tip and the outer feathers are white (Massam 2001). The white feathers can be seen most clearly when the bird is in flight. The bill, legs and feet are bright yellow, while the adult iris is reddish brown to brownish yellow in colour (Massam 2001). Male and female *A. tristis* are not clearly sexually dimorphic and are thus difficult to identify in the field (Counsilman Nee Jalil and Keng 1994).

Mynas are distinctive birds in that they move about in a walk rather than a hop. Like most territorial birds they have a bout of intense calling in the early morning that lasts between 5 and 15 minutes. Males call more often than the females, and pairs sometimes duet. The territorial call is a rowdy medley of creaky notes, growls, rattles, raucous, gurgling, chattering and bell-like sounds in rapid sequence often strung together as a song. Adults with young utter harsh squarking noises and young learning to fly emit persistent "chi-chi-chis". At their communal roosts mynas maintain a noisy chattering, even well after nightfall and before dawn. To hear samples of the common myna call please go to: [Tidemann, C. 2007b. Common Indian Myna Website > Identifying Mynas.](#)

Notes

Mynas in India are regarded as symbols of undying love, because they often pair for life. In India *maina* is used as a term of endearment for young girls (Tidemann 2007c).

The common myna is highly intelligent and acquires a fear response; it learns about an area in which it observes another individual experience an aversive event, namely capture by a human. This obviously has implications for management and control (Griffin & Boyce 2009).

Lifecycle Stages

The female myna incubates her eggs for 13 to 14 days. The fledging period lasts between 20 to 32 days, averaging 25 days. Parents feed the chicks as long as three weeks after they have left the nest (Massam 2001). Sexual maturity occurs at nine to 12 months. Juveniles form small flocks and may form mating pairs at as young as nine months old although few breed in their first year. Life span is an average of four years in the wild, possibly up to 12 years for some individuals (Markula Hannan-Jones & Csurhes 2009).

Uses

In India the common myna is referred to as the farmer's friend because it protects crops by feeding on insect pests. In fact the myna has been deliberately introduced to continental landmasses and islands with warm temperate to tropical climates ostensibly to control invertebrate pests (Case 1996, Veltman *et al.* 1996, Feare & Craig 1998).

Many myna species are accomplished mimics and can be taught to speak; for this reason the myna is a much sought-after pet in some parts of the world (Tidemann 2005), in Mallorca, Spain, several pet birds have escaped or been released into the wild.

Habitat Description

Common mynas are found in both tropical and temperate regions, from the tropics to southern Europe (Russia) and as far up as northern France (Feare 1998). They are able to adapt to a wide range of climates and habitats. They inhabit flood plains, grasslands, cultivated areas, plantations, as well as desert oases and foothills of various mountainous ranges (Feare and Craig, 1999). However, in general, the common myna reaches the highest densities in modified habitats near human establishments, including cities, towns, villages, farmland, rural dwellings, parks, gardens and roadsides (Gill 1999; Heather & Robertson 2000). The common myna evolved in open woodland habitats in India (Sengupta 1968, in Pell & Tidemann 1997) and is said to prefer anthropogenically modified woodland in Australia (Tidemann 2007e).

The common myna is good at adapting to local environments. For example, in Fiji it congregates on the seashore feeding off crustaceans and other stranded sealife and has even colonised a small coral island. In Singapore it is strongly associated with the rural landscape, for example, agricultural and farm areas (Lim Sodhi Brook and Soh 2003). In Australia mynas prefer reserve habitats, especially the perimeters (Pell and Tidemann 1997). While reserves provide excellent environs for the myna in Australia and stimulate large numbers of mynas during the breeding season, during the winter months mynas find refuge in the surrounding suburban areas (Pell and Tidemann 1997).

The common myna prefers warmer climates. For example, in New Zealand, it tends to avoid colder regions in the south such as Nelson; but interestingly it does establish stable populations near piggery sheds where sufficient heat is produced by the pigs to maintain a relatively high temperature; in addition there is an abundance of pig food available (P.R. Wilson Pers. Comm.). The common myna prefers to forage in open, grassy habitats (Crisp and Lill 2006, in Newey 2007), either in groups or alone, and roost in isolated stands of tall trees. In Singapore, it commonly roosts among monoclonal stands of tall densely canopied trees (Hails 1985; Yap *et al.* 2002). In Fiji, less densely canopied trees such as coconut palms are chosen by the adaptable bird for roosting and refuge (Stoner 1923).

Reproduction

The common myna stays in the same pair and maintains the same territory each year. It builds a cup-shaped nest out of dry grass, twigs and leaves and may construct its nest in a tree hollow, cliffside, building or thick vegetation. The common myna usually raises two broods per season, laying up to six pale greenish-blue eggs (3.1cm x 2.2cm; Massam 2001) in each brood, with an average of four. Both sexes brood and care for the young (Massam 2001). In New Zealand they lay eggs from mid-October to early March, with the highest egg-laying activity occurring from November to January. In Australia up to three broods of young may be produced in one season. Both sexes brood and care for the young (Massam 2001).

Nutrition

Common mynas are omnivorous scavengers and will feed on fruits, berries, grains, flower nectar, insects (including beetle larvae and adults, caterpillars, worms, flies, snails) and spiders. Nestlings are fed for the first ten days exclusively on invertebrates, primarily insects (Markula Hannan-Jones & Csurhes 2009). When insects are scarce, fruits and seeds make up a more important component of their diet (Peacock van Renburg & Robertson 2007). At such times, common mynas can become agricultural pests (Peacock van Renburg & Robertson 2007), feeding on the ripening fruit and seeds of plants such as figs, papaya, dates, apple, pear, tomato, and cereal crops such as maize, wheat and rice. Mynas are egg predators (Feare & Craig 1998), and are known to consume birds' eggs and chicks, as well as small reptiles.

Common mynas are highly adaptable to human habitations (Sontag & Louette 2007) and local food resources. For example, they have been known to consume pet food (Australian Museum 2003) and forage on the seashore for worms, molluscs, crustaceans and other seafood stranded on the mud flats (Stoner 1923). They also scavenge rubbish dumps, pastures, farmyards and roads for roadkill.

They are predominantly ground feeders, pecking prey from the surface in short pasture and grain stubble, but will opportunistically feed in flowering or fruiting trees and bushes (Feare and Craig 1999). In their native range of southern Asia the myna forms flocks in rural areas, which feast on insects and grubs turned up in the cultivated soil by the plough (Australian Museum 2003).

General Impacts

Flocks of the common myna are known to damage fruit crops, including grapes, apricots, apples, pears, strawberries, figs and gooseberries. (Heather and Robertson 1997).

Mynas are communal and commensal, they are highly vocal throughout the year, making them a public nuisance. Their droppings are a nuisance (Yap et al. 20002, in Lim Sodhi Brook and Soh 2003) and public health concern. Mynas form combined populations of up to 160 000 (Lim Sodhi Brook and Soh 2003) and roost in numbers as great as 5000 (Markula Hannan-Jones & Csurhes 2009). They are a residential nuisance as they build nests in spouting and drainpipes (Stoner 1923). Mynas fearlessly steal food off plates which may be a hygiene or general nuisance for restaurants and other shops and scavenge food from people's houses and gardens.

Common mynas pose a human health risk as they carry bird mites such as *Ornithonyssus bursa* and *Dermanyssus gallinae* that may infect humans. They can also cause dermatitis, asthma, severe irritation and rashes. Their droppings can spread Psittacosis, Ornithosis, Salmonellosis and arboviruses (Pers. comm. Bill Handke). They may also carry owl flies, biting lice, *Oxyspirrura* thread worm and round worm (Stoner 1923). Mynas are known to carry avian malaria (Massam 2001).

The common myna has been implicated in the demise of the lowland populations of the 'Vulnerable (VU)' Rarotonga starling (*Aplonis cinerascens*) (BirdLife International 2008b). Mynas are nest site competitors and can displace active breeding pairs of the Endangered (EN) Mauritius parakeet (*Psittacula eques*). In French Polynesia they are reported to predate on the Critically Endangered (CR) (*Todiramphus godeffroyi*).

Please follow this link for more examples of the [impacts of common mynas on threatened species](#).

Management Info

Preventative measures: Risk assessment models by the Bureau of Rural Sciences, Australia, classifies the common myna in the highest threat category (Bomford 2003). The common myna is prohibited in Western Australia (Massam 2001).

A Pest Animal Risk Assessment using a numerical risk assessment system developed by Bomford (2006) was carried out by the State of Queensland, Department of Employment, Economic Development and Innovation, in 2009. Indian mynas in Queensland were assessed as an 'extreme' threat species. See [Markula et al 2009](#)

Physical: Foraging traps are very useful for the control of small myna populations if poisoning is not an option. The Tindall Trap and the Tidemann Trap have been used successfully in New Zealand and Australia, respectively. The Decoy Trap, Kadavu Trap, Larsen Trap, Rat snap-trap and other foraging traps have also been used for trapping myna birds with less success. Please follow this link to view a [Trapping Mynas](#) and [Tidemann, C. 2007g. Common Indian Myna Website > Humane Disposal](#).

Chemical: Starlacide DRC1339 has been used against mynas and is effective where there are no non-target species issues. Alphachloralose paste is used for temporary local control of mynas in cooler climates. For more information on the use of these toxins please see [NZFSA. Undated. DRC 1339 For Bird Control](#) and Nelson. 1994. *Bird Control in New Zealand Using Alpha-Chloralose and DRC1339*.

Integrated Pest Management: As invasive bird species are frequently associated with human modified environments IPM is an appropriate strategy (Lim Sodhi Brook and Soh 2003). Long term management practices may include habitat modification, resource limitation and public education. Restricting food available to the myna is difficult as it has such a variable diet (Thomas 2004).

The need to raise public awareness is important part of IPM. Envirotalk Australia has a [forum discussion](#) on the myna topic. The [Minimising Myna Website](#) aids public education on the issue. The [Canberra Indian Myna Action Group](#) is a community group that has developed a number of strategies, including public education and a trapping program, to tackle the common myna. CIMAG's trapping program has been highly successful and has humanely removed over 12 000 Mynas from around Canberra in around 18 months (CIMAG Undated).

Research: Kate Garrock is undertaking a PhD research project at Australian National University on the impact of Indian Mynas on native birds and the impact of trapping activity in the Canberra region.

Pathway

In Israel, mynas escaped from a private facility of exotic birds in the centre of the Tel Aviv public park. On oceanic islands, invasion pathways appear to be primarily via ships, particularly large ferries (Tearika 2003, D. Wattling Pers. Comm.). The pathway to the Spanish islands has been through pet shops and later escapes from the home cages. Introduced by acclimatisation societies.

Principal source:

Compiler: IUCN/SSC Invasive Species Specialist Group (ISSG)

Updates on management information with support from the Overseas Territories Environmental Programme (OTEP) project XOT603, a joint project with the Cayman Islands Government - Department of Environment

Review:

Publication date: 2011-07-04

ALIEN RANGE

[1] AMERICAN SAMOA
[2] BRUNEI DARUSSALAM
[8] COOK ISLANDS
[1] FRANCE
[1] HONG KONG
[1] INDONESIA

[10] AUSTRALIA
[1] COMOROS
[4] FIJI
[19] FRENCH POLYNESIA
[2] INDIA
[1] IRAQ

[2] ISRAEL
[1] KUWAIT
[1] MALDIVES
[1] MAYOTTE
[5] NEW ZEALAND
[1] QATAR
[1] RUSSIAN FEDERATION
[2] SAMOA
[6] SEYCHELLES
[4] SOLOMON ISLANDS
[1] SPAIN
[1] TURKEY
[8] UNITED STATES
[2] VANUATU

[2] KIRIBATI
[1] MADAGASCAR
[1] MAURITIUS
[2] NEW CALEDONIA
[1] OMAN
[1] REUNION
[3] SAINT HELENA
[1] SAUDI ARABIA
[1] SINGAPORE
[1] SOUTH AFRICA
[1] TONGA
[1] UNITED ARAB EMIRATES
[2] UNITED STATES MINOR OUTLYING ISLANDS
[1] WALLIS AND FUTUNA

Red List assessed species 59: CR = 5; EN = 7; VU = 11; NT = 4; LC = 32;

[Acanthiza reguloides](#) LC
[Acrocephalus caffer](#) EN
[Acrocephalus mendanae](#) LC
[Acrocephalus sechellensis](#) NT
[Cacatua galerita](#) LC
[Callocephalon fimbriatum](#) VU
[Collocalia leucophaea](#) VU
[Copsychus sechellarum](#) EN
[Cormobates leucophaea](#) LC
[Cyanoramphus novaezelandiae](#) VU
[Dendrocopos syriacus](#) LC
[Grallina cyanoleuca](#) LC
[Humblotia flavirostris](#) EN
[Jynx ruficollis](#) LC
[Onychoprion fuscatus](#) LC
[Otus insularis](#) EN
[Pachycephala rufiventris](#) LC
[Passer melanurus](#) LC
[Philesturnus carunculatus](#) NT
[Platycercus eximius](#) LC
[Polytelis swainsonii](#) VU
[Pomarea nigra](#) CR
[Psittacula eque](#) EN
[Puffinus pacificus](#) LC
[Rhipidura albiscapa](#) LC
[Rhipidura leucophrys](#) LC
[Terpsiphone corvina](#) VU
[Todiramphus ruficollaris](#) VU
[Trachyphonus vaillantii](#) LC
[Zosterops modestus](#) EN

[Acanthorhynchus tenuirostris](#) LC
[Acrocephalus kerearako](#) NT
[Acrocephalus rimatarae](#) VU
[Aplonis cinerascens](#) VU
[Cacatua roseicapilla](#) LC
[Charadrius sanctaehelenae](#) CR
[Copsychus saularis](#) LC
[Coracina typica](#) VU
[Cracticus torquatus](#) LC
[Dacelo novaeguineae](#) LC
[Gerygone igata](#) LC
[Hemiphaga novaeseelandiae](#) NT
[Hypsipetes olivaceus](#) VU
[Malurus cyaneus](#) LC
[Otus capnodes](#) CR
[Otus pauliani](#) CR
[Pardalotus striatus](#) LC
[Petaurus breviceps](#) LC
[Platycercus elegans](#) LC
[Ploceus velatus](#) LC
[Pomarea mendozae](#) EN
[Prothemadara novaeseelandiae](#) LC
[Ptilinopus rarotongensis](#) VU
[Pyrrholaemus sagittatus](#) LC
[Rhipidura fuliginosa](#) LC
[Stigmatopelia senegalensis](#) LC
[Todiramphus godeffroyi](#) CR
[Todiramphus sanctus](#) LC
[Zosterops lateralis](#) LC

BIBLIOGRAPHY

129 references found for *Acridotheres tristis*

Management information

ACT 2003. State of the Environment Report 2003. Indicator-Pest Animals. Australian Capital Territory.

Atkinson, I. A. E. and Atkinson, T. J. 2000. Land vertebrates as invasive species on islands served by the South Pacific Regional Environment Programme. In: *Invasive Species in the Pacific: A Technical Review and Draft Regional Strategy*. South Pacific Regional Environment Programme, Samoa: 19-84.

Summary: This report reviews available information on the adverse effects of 14 alien vertebrates considered to be significant invasive species on islands of the South Pacific and Hawaii, supplementing the authors' experience with that of other workers.

Bomford, M., 2003. *Risk Assessment for the Import and Keeping of Exotic Vertebrates in Australia*. Bureau of Rural Sciences, Canberra.

Summary: Available from: <http://www.feral.org.au/wp-content/uploads/2010/03/PC12803.pdf> [Accessed August 19 2010]

Canning, Gregory, 2011. *Eradication of the invasive common myna, *Acridotheres tristis*, from Fregate Island, Seychelles*. *Phelsuma* 19 (2011); 43-53

Summary: Available from: <http://www.islandbiodiversity.com/Phelsuma%2019d.pdf> [Accessed 2 July 2011]

Department of Agriculture and Food, 2008. *National Animal Pest Alert: Common myna*. Department of Agriculture and Food the State of Western Australia

Summary: Available from: http://www.agric.wa.gov.au/content/PW/vp/bird/CommonMyna_NHT.pdf [Accessed 3 July 2008]

Doherty, N. 2006. *Information booklet for the "National Control of the myna" (*Acridotheres spp.*)*. Government of Samoa Division of Environment & Conservation

Hails, C. J. 1985. Studies of problem bird species in Singapore: I. Sturnidae (Mynas and Starlings). A report submitted to the commissioner for Parks and Recreation, Ministry of National Development, Singapore.

Summary: Notes on invasive characteristics.

Hatzofe, O. and Perelman, Y. 2001. Myna trapping trial report: summary and recommendations. Israel Nature and Parks Authority internal report of the Science and Conservation Division. 4pp.

Summary: In Hebrew.

Hughes, B.J., G.R. Martin & S.J. Reynolds. 2008. Cats and seabirds: effects of feral Domestic Cat *Felis silvestris catus* eradication on the population of Sooty Terns *Onychoprion fuscatus* on Ascension, *Ibis* 150(Suppl. 1): 122-131.

IUCN 2010. *IUCN Red List of Threatened Species*. Version 2010.4.

Summary: The IUCN Red List of Threatened Species provides taxonomic, conservation status and distribution information on taxa that have been globally evaluated using the IUCN Red List Categories and Criteria. This system is designed to determine the relative risk of extinction, and the main purpose of the IUCN Red List is to catalogue and highlight those taxa that are facing a higher risk of global extinction (i.e. those listed as Critically Endangered, Endangered and Vulnerable). The IUCN Red List also includes information on taxa that are categorized as Extinct or Extinct in the Wild; on taxa that cannot be evaluated because of insufficient information (i.e. are Data Deficient); and on taxa that are either close to meeting the threatened thresholds or that would be threatened were it not for an ongoing taxon-specific conservation programme (i.e. are Near Threatened).

Available from: <http://www.iucnredlist.org/> [Accessed 25 May 2011]

IUCN/SSC Invasive Species Specialist Group (ISSG), 2010. *A Compilation of Information Sources for Conservation Managers*.

Summary: This compilation of information sources can be sorted on keywords for example: Baits & Lures, Non Target Species, Eradication, Monitoring, Risk Assessment, Weeds, Herbicides etc. This compilation is at present in Excel format, this will be web-enabled as a searchable database shortly. This version of the database has been developed by the IUCN SSC ISSG as part of an Overseas Territories Environmental Programme funded project XOT603 in partnership with the Cayman Islands Government - Department of Environment. The compilation is a work under progress, the ISSG will manage, maintain and enhance the database with current and newly published information, reports, journal articles etc.

Kang, N., Sigurdsson, J. B., Hails, C. J., & Counsilman J. J. 1990. Some implications of resource removal in the control of mynas (*Acridotheres spp.*) in Singapore. *Malayan Nature Journal* 44: 103-108.

Summary: Long term management issues.

Lim, H.C., Sodhi, N.S., Brook, B.W. and Soh, M.C.K. 2003. Undesirable Aliens: Factors Determining the Distribution of Three Invasive Bird Species in Singapore, *Journal of Tropical Ecology* 19: 685-695.

Summary: Land use in Singapore and the presence of invasive birds.

Live Arico. 2007. First Campaign for Common Myna (*Acridotheres tristis*) Control in the island of Mallorca 2006.

Live Arico Tenerife Animal and Environmental Protection, 2007. First campaign for Common Myna (*Acridotheres tristis*) control in the island of Mallorca. Final Report for the Environment Department Balearic Islands' Government

Markula, A., Hannan-Jones, M. & Csorhes, S., (2009). *Pest animal risk assessment: Indian myna (*Acridotheres tristis*)*. *Biosecurity Queensland. Queensland Primary Industries and Fisheries*.

Summary: Available from: http://www.dpi.qld.gov.au/documents/Biosecurity_EnvironmentalPests/IPA-Indian-Myna-Risk-Assessment.pdf [Accessed 30 October 2009]

Millett, J, Climo, G & Jivan Shah, N. 2004. Eradication of the common myna *Acridotheres tristis* populations in the granitic Seychelles: successes, failures and lessons learned. *Advances in Vertebrate Pest Management* 3: 169-183.

Nelson, P.C. 1994. Bird Control in New Zealand Using Alpha-Chloralose and DRC1339, *Vertebrate Pest Conference Proceedings collection: Proceedings of the Sixteenth Vertebrate Pest Conference*.

Northland Regional Council. Undated. [Home > Environment > Weed and pest control > Pest animals > Myna](#)

Summary: Available from: <http://www.nrc.govt.nz/Environment/Weed-and-pest-control/Animal-pests/Myna/> [Accessed 27 October 2009]
 NZFSA (New Zealand Food Safety Authority). Undated. [home > acvm > publications > notes > Drc1339 Bird Study Notes: Controlled Pesticides: DRC 1339 For Bird Control](#).

Summary: Available from: <http://www.nzfsa.govt.nz/acvm/publications/notes/drc1339-bird-study-notes.pdf> [Accessed 30 October 2009]
 Pacific Invasives Initiative (PII). Undated. *A major myna matter: managing invasive birds in the Pacific*.

Parkes, J. 2009. Common Mynas (*Acridotheres tristis*) on the Three Kings Islands: Should and Can They Be Controlled or Eradicated?
 Pierce, R.J. 2005. *A Preliminary Review of Interactions Between Introduced Mynas and Indigenous Vertebrate Fauna and Methods for Controlling Mynas*.

RSPCA (2017). *Management of common (Indian) myna birds*, July 2017

[Saavedra, Susana, 2009. First control campaign for common myna *Acridotheres tristis* on Ascension Island. Live Arico Invasive Species Department](#)

Savage, J. (Bird Gard). 2004. Subject: Re: [Aliens-L] Dealing with Common Mynas (*Acridotheres tristis*). Email communication.

Tidemann, C. (2005). Indian Mynas - Can the problems be controlled? In Urban Animal Management Conference Proceedings 2005. The Australian National University, pp. 55-57.

[Tidemann, C. 2007a. Common Indian Myna Website > Home. The Australian National University.](#)

Summary: Useful information, including how to trap and humanely dispose of mynas. Limited stocks of myna traps are available for sale - if you wish to purchase a trap please express interest via the web site. If you represent a group and would like your web site to be linked to the anu myna site - please email myna@anu.edu.au with Myna Group in the subject line and the url of your web site in the text of the message. Please keep your message brief.

Available from: <http://fennerschool-associated.anu.edu.au/myna/index.html> [Accessed 27 October 2009]

[Tidemann, C. 2007e. Common Indian Myna Website > Minimising Mynas: Mitigation of the impact of Mynas on biodiversity and public amenity. The Australian National University.](#)

Summary: The web site provides information on a new, expanded, trial of myna traps in Canberra suburbs commencing in March 2004.

Available from: <http://fennerschool-associated.anu.edu.au/myna/minimising.html> [Accessed 27 October 2009]

[Tidemann, C. 2007f. Common Indian Myna Website > Trapping Mynas: Humane bird trap for Common Indian Mynas and European Starlings. The Australian National University.](#)

Summary: Available from: <http://fennerschool-associated.anu.edu.au/myna/trapping.html> [Accessed 27 October 2009]

[Tidemann, C. 2007g. Common Indian Myna Website > Humane Disposal. The Australian National University.](#)

Summary: Available from: <http://fennerschool-associated.anu.edu.au/myna/humane.html> [Accessed 27 October 2009]

[Tidemann, C. 2007h. Common Indian Myna Website > Monitoring Mynas. The Australian National University.](#)

Summary: Available from: <http://fennerschool-associated.anu.edu.au/myna/monitoring.html> [Accessed 27 October 2009]

Tidemann, C.R. (2001). Mitigation of the impact of mynas on biodiversity and public amenity. The Australian National University. School of Resources, Environment & Society.

Tindall, S.D., C.J. Ralph & M.N. Clout. 2007. Changes in bird abundance following common myna control on a New Zealand island, *Pacific Conservation Biology* 13(3): 202-212.

[Varnham, K. 2006. Non-native species in UK Overseas Territories: a review. JNCC Report 372. Peterborough: United Kingdom.](#)

Summary: This database compiles information on alien species from British Overseas Territories.

Available from: <http://www.jncc.gov.uk/page-3660> [Accessed 10 November 2009]

[Wildland Consultants. 2000. Norske Skog Tasman Wildlands: Wetland Restoration Project.](#)

Summary: Integrated management of myna by the restoration of vulnerable island ecosystems.

Available from: http://www.wildland.co.nz/projects/norsketasman/norske_project.htm [Accessed 07 December 2005]

Burns, F., McCulloch, N., Székely, T. & Bolton, M. (2013). The impact of introduced predators on an island endemic, the St Helena Plover, *Charadrius sanctaehelenae*. *Bird Conserv. Int.* 23, 125-135

Charter, M., Izhaki, I., Ben Mocha, Y. & Kark, S. (2016). Nest-site competition between invasive and native cavity nesting birds and its implication for conservation. *J. Environ. Manage.* 181, 129-134

Feare, C.J., Lebarbenchon, C., Dietrich Larose, S. (2015). Predation of seabird eggs by Common Mynas *Acridotheres tristis* on Bird Island, Seychelles, and its broader implications. *Bull ABC* 22:162-170

Grarock, K. et al., (2014). Are invasive species drivers of native species decline or passengers of habitat modification? A case study of the impact of the common myna (*Acridotheres tristis*) on Australian bird species. *Austral Ecology*, 39(1), pp.106-114.

Grarock, K., Lindenmayer, D.B., Wood, J.T., and Tidemann, C.R. (2013). Does human-induced habitat modification influence the impact of introduced species? A case study on cavity-nesting by the introduced Common Myna (*Acridotheres tristis*) and two Australian native parrots. *Environmental Management* 52, 958-970.

Grarock, K., Tidemann CRW, J., Lindenmayer, D. (2012). Is it benign or is it a pariah? Empirical evidence for the impact of the common myna (*Acridotheres tristis*) on Australian birds. *PLoS ONE* 7:e40622

Haythorpe, K. (2013). Competitive behaviour in Common Mynas. An investigation into potential impacts on native fauna and solutions for management. Thesis submitted to Charles Sturt University in partial fulfilment of the requirements of the Doctor of Philosophy (PhD).

Available from: <https://researchoutput.csu.edu.au/en/publications/competitive-behaviour-in-common-mynas-an-investigation-into-poten-3>. (Accessed: 31st March 2020).

Hughes, B.J., Dickey, R.C., & Reynolds, S.J. (2019). Predation pressures on sooty terns by cats, rats and common mynas on Ascension Island in the South Atlantic. In C. R. Veitch, M. N. Clout, A. R. Martin, J. C. Russell, & C. J. West (Eds.), *Island Invasives: Scaling up to Meet the Challenge* (pp. 291-297). Gland, Switzerland: IUCN

Hughes, B. J., Martin, G. R. & Reynolds, S. J. (2017). Estimating the extent of seabird egg depredation by introduced Common Mynas on Ascension Island in the South Atlantic. *Biol. Invasions* 19, 843-857

Lowe, K.A., Taylor, C.E., and Major, R.E. (2011). Do common mynas significantly compete with native birds in urban environments? *Journal of Ornithology* 152, 909-921.

Pell, A.S., and Tidemann, C.R. (1997). The impact of two exotic hollow-nesting birds on two native parrots in savannah and woodland in eastern Australia. *Biological Conservation* 79, 145-153.

Thibault, J. C., Martin, J. L., Penloup, A., and Meyer, J. Y. (2002). Understanding the decline and extinction of monarchs (Aves) in Polynesian islands. *Biological Conservation* 108, 161-174. doi:10.1016/S0006-3207(02)00102-7

General information

Armstrong D.P.; Ewen J.G.; Dimond W.J.; Lovegrove T.G.; Bergstrom A.; Walter B. 2000. Breeding biology of North Island robins (*Petroica australis longipes*) on Tiritiri Matangi Island, Hauraki Gulf, New Zealand. *Notornis* 47: 106-118

[Australian Museum. 2003. Common Myna *Acridotheres tristis*.](#)

Summary: Biological information.

Available from: <http://www.birdsinbackyards.net/finder/display.cfm?id=36> [Accessed 10 November 2009]

[Avibase, undated. The world bird database: Common Myna \(*Acridotheres tristis*\) \(Linnaeus, 1766\)](#)

Summary: Available from:

<http://www.bsc-eoc.org/avibase/species.jsp?lang=EN&id=D1CF6D3C0FE5774D&ts=1203996833593&sec=summary> [Accessed 27 February 2008]

Beichle, U. R. 1989. Common myna on Upolu: first record for the Western Samoa Islands. *Elepaio* 49: 85-86.

Summary: Distribution.

Blanvillain, Caroline & Salducci, J. & Tutururai, G & Maeura, M. (2003). Impact of introduced birds on the recovery of the Tahiti Flycatcher (*Pomarea nigra*), a critically endangered forest bird of Tahiti. *Biological Conservation - BIOL CONSERV.* 109. 197-205.
10.1016/S0006-3207(02)00147-7.

Blanvillain C.; Pierce R.; Saducchi J-M. 1999. Trimestrial report on the reproductive success of the O'mamao or Tahiti flycatcher (*Pomarea nigra*). Unpublished report to SPREP

Bull, P.C.; Gaze, P.D.; Robertson, C.J.R. 1985. The Atlas of Bird Distribution in New Zealand. Ornithological Society of New Zealand.

Burt, April & Gane, Julie & Olivier, Iain & Calabrese, Licia & GROENE, ARJAN & LIEBRICK, TANYA & MARX, DANE & Shah, Nirmal. (2016). The history, status and trends of the Endangered Seychelles Magpie-robin *Copsychus sechellarum*. *Bird Conservation International*. -1. 1-19.
10.1017/S0959270915000404.

Byrd, G. V. 1979. Common myna predation on wedge-tailed shearwater eggs. *Elepaio* 39: 69-70.

Summary: Impacts.

Byrd, G. V., Moriarty, D. I. & Brady, B. G. (1983). Breeding biology of Wedge-tailed Shearwaters at Kilauea Point, Hawaii. *Condor* 85, 292-296
CABI. (2009). Invasive Species Compendium. Datasheet report for *Acridotheres tristis* (common myna).

[CIMAG. Undated. Indian myna: the Flying Cane Toad.](#)

Summary: Available from: <http://www.indianmynaaction.org.au/documents/Brochure%20-%20the%20Flying%20Cane%20Toad%201.pdf> [Accessed 30 October 2009]

[Cook Islands Biodiversity Database. Undated. *Acridotheres tristis* Manu Kavamani, Common Myna. Cook Islands Biodiversity & Natural Heritage.](#)

Summary: Available from: <http://cookislands.bishopmuseum.org/species.asp?id=8262> [Accessed 29 October 2007]

Counsillman, J.J., Nee, K., Jalil, A.K. and Keng, W.L. 1994. Discriminant Analysis of Morphometric Characters as a Means of Sexing Mynas, *Journal of Field Ornithology*, Vol. 65(1): 17.

Summary: In depth study of physical differences between male and female mynas.

Crisp, H., Lill, A. (2006). City slickers: habitat use and foraging in urban common mynas *Acridotheres tristis*. *Corella* 30:9-15

CSN 1972. Classified Summarised Notes. *Notornis* 23: 89.

CSN 1974. Classified Summarised Notes. *Notornis* 25.

CSN 1981. Classified Summarised Notes. *Notornis* 28: 77.

CSN 1984. Classified Summarised Notes. *Notornis* 31: 55.

CSN 1994. Classified Summarised Notes, North Island. *Notornis* 41: 49.

Department of Environment and Climate Change (NSW), (2007). Terrestrial Vertebrate Fauna of the Greater Southern Sydney Region: Volume 2 Fauna of Conservation Concern & Priority Pest Species. A joint project between the Sydney Catchment Authority and the Department of Environment and Climate Change (NSW) (DECC).

[Dhami, M.K. & Nagle, B. 2009. Review of the Biology and Ecology of the Common Myna \(*Acridotheres tristis*\) and some implications for management of this invasive species. PII \(Pacific Invasives Initiative\).](#)

Summary: Available from: http://www.issg.org/cii/Electronic%20references/pii/Myna_review_final.pdf [Accessed 10 November 2009]

Drent R. 1996. Myna eradication boosts *tuis*. *Sunday Star Times* 4, August 1996.

Eguchi, K. & Amano, H. E. (2004). Invasive birds in Japan. *Global Environmental Research* 8: 29-39.

Fares, K., Muna, S. and Mohammed, A. (2021). Anthropogenic not climatic correlates are the main drivers of expansion of non-native common myna *Acridotheres tristis* in Jordan. *Management of Biological Invasions* (2021) Volume 12, Issue 3: 640-653

Feare, C., and A. Craig. 1999. Starlings and mynas. Illustrated by Barry Croucher, Chris Shields and Kamol Komolphalin. Princeton University Press Princeton New Jersey

Summary: Description, habitat.

Feare, C. & Craig, A. 1998. Starlings and mynas. Christopher Helm (A & C Black), London, United Kingdom. 285 pp.

Summary: Notes on uses.

Fleischer, R.C., R.N. Williams & A.J. Baker. 1991. Genetic Variation within and among Populations of the Common Myna (*Acridotheres tristis*) in Hawaii, *Journal of Heredity* 82: 205-208.

Foster, J.T. & S.K. Robinson. 2007. Introduced Birds and the Fate of Hawaiian Rainforests, *Conservation Biology* 21(5): 1248-1257.

Freifeld, H.B. (1999) Habitat relationships of forest birds on Tutuila Island, American Samoa. *Journal of Biogeography* 26, 1191-1213.

[Gargominy, O., Bouchet, P., Pascal, M., Jaffre, T. and Tourmeu, J. C. 1996. Consequences des introductions d'especes animales et vegetales sur la biodiversite en Nouvelle-Caledonie. Rev. Ecol. \(Terre Vie\) 51: 375-401.](#)

Summary: Consequences to the biodiversity of New Caledonia of the introduction of plant and animal species.

Gibson-Hill, C. A. 1949. An annotated checklist of the birds of Malaya. *Bulletin of Raffles Museum* 20:1-299.

Summary: Description of myna.

Gill, B. J. 1999. A myna increase - notes on introduced mynas (*Acridotheres*) and bulbuls (*Pycnonotus*) in Western Samoa. *Notornis* 46: 268-269.

Summary: Invasive characteristics.

[Government of Western Australia Undated. National Pest Animal Alert: Common Myna. Government of Western Australia Department of Food and Agriculture.](#)

Summary: Fact Sheet

Available from: http://www.agric.wa.gov.au/objtwr/imported_assets/content/pw/vp/bird/commonmyna_nht.pdf [Accessed 04 November 2009]

Griffin, A.S. 2009. Temporal Limitations on Social Learning of Novel Predators by Indian Mynahs, *Acridotheres tristis*, *Ethology* 115:287-295.—

Global Invasive Species Database (GISD) 2025. Species profile *Acridotheres tristis*. Available from: <https://iucngisd.org/gisd/species.php?sc=108> [Accessed 30 July 2025]

Griffin, A.S. & H.M. Boyce. 2009. Indian mynahs, *Acridotheres tristis*, learn about dangerous places by observing the fate of others, *Animal Behaviour* 78: 79-84.

Handal, E. & M.B. Qumsiyeh. (2021). Status and distribution of the invasive Common Myna *Acridotheres tristis* in the West Bank, Palestine. *Sandgrouse*, 43:129-132

Harper, M. J., McCarthy, M. A. & Van Der Ree, R. (2005). The use of nest boxes in urban natural vegetation remnants by vertebrate fauna. *Wildl. Res.* 32, 509-516

Hatzofe, O. 2001. Pers. Comm. Re: *FW: Indian Myna in West Africa* (Email communication.)

Summary: Overview of control measures in place in Israel against the common myna c. 2001.

Heather B.D.; Robertson H.A. 2000. The new field guide to the birds of New Zealand. Viking, Auckland.

Heptonstall, R. E. A. (2010). The distribution and abundance of Myna Birds (*Acridotheres tristis*) and Rimatara Lorikeets (*Vini kuhlii*) on Atiu, Cook Islands. MSc Thesis. Submitted in accordance with the requirements for the degree of Masters of Science.

Holyoak, D. T. and Thibault, J. C. 1984. Contribution a l'Étude des Oiseaux de Polynésie Orientale. Mémoires du Muséum National d'Histoire Naturelle, n.s. Série A, Zoologie, Tome 127. Éditions du Museum, Paris.

Holzapfel, C., Levin, N., Hatzofe, O. & Kark, S. (2006). Colonisation of the Middle East by the invasive Common Myna *Acridotheres tristis* L., with special reference to Israel. *Sandgrouse* 28, 44-51

[ITIS \(Integrated Taxonomic Information System\)](http://www.itis.gov/), 2004. Online Database *Acridotheres tristis*

Summary: An online database that provides taxonomic information, common names, synonyms and geographical jurisdiction of a species. In addition links are provided to retrieve biological records and collection information from the Global Biodiversity Information Facility (GBIF) Data Portal and bioscience articles from BioOne journals.

Available from: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=554025 [Accessed December 31 2004]

Kang, N. 1989. Comparative behavioural ecology of the mynas, *Acridotheres tristis* (Linnaeus) and *A. javanicus* (Cabanis) in Singapore. PhD thesis, National University of Singapore.

Summary: Behaviour and ecology.

Komdeur, J. (2008). Breeding of the Seychelles Magpie Robin *Copsychus sechellarum* and implications for its conservation. *Ibis*, 138, 485-498.

Landcare Research (LCR). 2008. Garden Bird Survey. Landcare Research, New Zealand.

Lever C. 1987. Naturalized Birds of the World. Edition Longman Scientific & Technical, Avon, U.K. 615 pp.

Lever, C. 1994. Naturalized Animals: the Ecology of Successfully Introduced Species. London: T. & A. D. Poyer.

Lieberman, A., McCormack, G. (2008). An island endemic: reintroduction of the Kuhl's Lory. *PsittaScene* 20(4): 8-11.

Live Arico. Undated. *Miná común - Common myna*.

[Martin, W.K. 1996. The Current and Potential Distribution of the Common Myna *Acridotheres tristis* in Australia, *Emu* 96: 166-173.](http://www.publish.csiro.au/?act=view_file&file_id=MU9960166.pdf)

Summary: Available from: http://www.publish.csiro.au/?act=view_file&file_id=MU9960166.pdf [Accessed March 2 2010]

Massam, M. 2001. *Common myna* [Farmnote No. 61/2001]. Department of Agriculture (Western Australia).

McAllan, I.A.W. & D. Hobcroft. 2005. The further spread of introduced birds in Samoa, *Notornis* 52: 16-20.

Mc Kenzie H.R. 1979. A history and account of the birds of the Hunua Ranges. *Notornis* 26: 105-119.

Monson, V. 2005. Bio-control the Last Resort for Hawaii's Plants, *The Maui News*.

Summary: Mentions the role of the common myna on the spread of invasive plants in Hawaii (which are a major threat to the survival of Hawaii's native flora).

[Muséum national d'Histoire naturelle \[Ed\]. 2003-2006. *Acridotheres tristis*. Inventaire national du Patrimoine naturel, site Web : http://inpn.mnhn.fr. Document téléchargé le 28 mars 2008.](http://inpn.mnhn.fr/Document_téléchargé_le_28_mars_2008)

Summary: Available

from: http://inpn.mnhn.fr/isb/servlet/ISBServlet?action=Espece&typeAction=10&pageReturn=ficheEspeceDescription.jsp&numero_taxon=199444 [Accessed March 20 2008]

Newey, Philip. 2007. Foraging behaviour of the Common Myna (*Acridotheres tristis*) in relation to vigilance and group size, *Emu* 107: 315-320.

Newsome, A.E. & Noble, I.R. 1986. Ecological and physiological characters of invading species. Pp. 1-20 in *Ecology of Biological Invasions: An Australian Perspective*. Eds R.H. Groves & J.J. Burdon. Australian Academy of Science, Canberra.

[Orchan Y, Chiron F, Schwartz A, Kark S \(2013\) The complex interaction network among multiple invasive bird species in a cavity-nesting community. *Biological Invasions* 15: 429-445. https://doi.org/10.1007/s10530-012-0298-6](https://doi.org/10.1007/s10530-012-0298-6)

Parsons, H., Major, R.E., French, K. (2006). Species interactions and habitat associations of birds inhabiting urban areas of Sydney, Australia. *Austral Ecol* 31:217-227

Pascal, M., Barré, N., De Garine-Wichatitsky, Lorvelec, O., Frétey, T., Brescia, F., Jourdan, H. 2006. Les peuplements néo-calédoniens de vertébrés : invasions, disparitions. Pp 111-162, in M.-L. Beauvais et al., : *Les espèces envahissantes dans l'archipel néo-calédonien*, Paris, IRD Éditions, 260 p.+ cd-rom

Summary: Synthèse des introductions d'espèces de vertébrés en Nouvelle-Calédonie et évaluation de leurs impacts.

Peacock, D.S., van Renburg, B.J., Robertson, M.P. 2007. The distribution and spread of the invasive alien common myna, *Acridotheres tristis* L. (Aves: Sturnidae), in southern Africa, *South African Journal of Science* 103(11-12): pp. 465-473.

[Pell, A. S., & Tidemann, C. R. \(1997\). The Ecology of the Common Myna in Urban Nature Reserves in the Australian Capital Territory. *Emu - Austral Ornithology*, 97\(2\), 141-149. https://doi.org/10.1071/MU97018](https://doi.org/10.1071/MU97018)

Summary: Mynas and their association with reserves.

Pernetta, J. C. and Watling, D. 1978. The introduced and native terrestrial vertebrates of Fiji. *Pacific Science* 32: 223-244.

Pierce R.J.; Blanvillain C. 2000. Conservation of the Tahiti flycatcher (*Pomarea nigra*): report on advice provided to Societe d'Ornitologie de Polynesie. Conservation Advisory Science Notes 290. Department of Conservation, Wellington. 8 pp.

Probst J.-M. 1997. Animaux de la Réunion. Azalées Éditions. 168 pp.

Rajan, P. & Pramod, P. (2013). Introduced birds of the Andaman & Nicobar Islands, India. *Indian Birds*, 8(3), pp.71-72.

[Science Centre. Undated. *Faunal Wealth of Andaman & Nicobar Islands*.](http://www.and.nic.in/Know%20Andaman/faunalwealth/faunal_wealth.htm)

Summary: Available from: http://www.and.nic.in/Know%20Andaman/faunalwealth/faunal_wealth.htm [Accessed 03 November 2009]

Seitre, R. & Seitre, J. 1992. Causes of land-bird extinction in French Polynesia. *Oryx*, 26, 215-222.

Summary: Les auteurs ont rassemblé des données écologiques sur les oiseaux terrestres endémiques, les animaux introduits et les habitats. Avec la chasse et la destruction des habitats, les introductions de prédateurs ont joué un rôle majeur dans la régression et l'extinction de plusieurs espèces d'oiseaux. L'éradication rapide de ces prédateurs, associée à des re-introductions d'oiseaux, constituerait la manière la plus efficace pour assurer la survie des espèces restantes.

Stoner, D. 1923. The Mynah. A Study in Adaptation, *Auk* 40(2): 328-330.

Summary: Comments on myna impacts in Fiji.

[The Cook Islands Natural Heritage Trust. 2005. *Acridotheres tristis*: Manu Kavamani \(Common Myna\).](#)

Summary: Cook Islands Biodiversity database

Available from: <http://cookislands.bishopmuseum.org/search.asp> [Accessed 08 December 2005]

Thibault, J.C., 1998. Extrait du compte-rendu de la seconde mission. *Te Manu* 25, 2.

Thomas, A. 2004. Myna Fightback, In *Scribbly Gum*, ABC. 2005.

Summary: Article on trapping mynas.

[Tidemann, C. 2005. *Common Myna*. The Australian National University.](#)

Summary: Biological information.

Available from: <http://sres.anu.edu.au/associated/myna/problem.html> [Accessed 06 December 2005]

[Tidemann, C. 2007b. *Common Indian Myna Website > Identifying Mynas*. The Australian National University.](#)

Summary: Available from: <http://fennerschool-associated.anu.edu.au//myna/identification.html> [Accessed 27 October 2009]

[Tidemann, C. 2007c. *Common Indian Myna Website > Myna Problems*. The Australian National University.](#)

Summary: Available from: <http://fennerschool-associated.anu.edu.au//myna/problem.html> [Accessed 27 October 2009]

[Tidemann, C. 2007d. *Common Indian Myna Website > Are Mynas spreading?*. The Australian National University.](#)

Summary: Available from: <http://fennerschool-associated.anu.edu.au//myna/spreading.html> [Accessed 27 October 2009]

U.S. Fish and Wildlife Service. (2005). Regional Seabird Conservation Plan, Pacific Region. U.S. Fish and Wildlife Service, Migratory Birds and Habitat Programs, Pacific Region, Portland, Oregon

van Rensburg, B.J., D.S. Peacock & M.P. Robertson. 2009. Biotic homogenization and alien bird species along an urban gradient in South Africa, *Landscape and Urban Planning*.

Yap, C. A. M., N. S. Sodhi & B. W. Brook. 2002. Roost characteristics of invasive mynas in Singapore. *Journal of Wildlife Management* 66: 1118-1127.

Summary: Roost characteristics.