

FULL ACCOUNT FOR: Cecropia peltata





System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Plantae	Magnoliophyta	Magnoliopsida	Urticales	Cecropiaceae

snakewood tree (English), yagrumo hembra (Spanish), faux ricin (French), Common name

pisse-roux (French), bois cannon (French), pop-a-gun (English), trumpet wood (English), guarumo (Spanish), Trompetenbaum (German), parasolier (French),

trumpet tree (English), papyrus géant (English)

Synonym Ambaiba pelata , Kuntze

Coilotapalus peltata, Britton

Cecropia schreberiana Similar species

Cecropia peltata is a fast-growing, short-lived tree that grows in neotropical **Summary**

regions. It is light-demanding and rapidly invades disturbed areas, such as forest canopy gaps, roadsides, lava flows, agricultural sites, urban locations, and other disturbed areas. It naturally occurs in tropical Central and South America, as well as some Caribbean islands and has been introduced to Malaysia, Africa, and Pacific Islands. It may be replacing, or competing with,

other native pioneer species in some locations.



view this species on IUCN Red List

Species Description

Cecropia peltata is a neotropical tree that reaches heights of 20 m or more. Its stems are hollow, partitioned at the nodes, and bear U-shaped leaf scars. Its leaves are alternate, long-lobed, ovate, somewhat pointed, about 10-50 cm wide, dark green and scaborous above and densley white-tomentose underneath. Its staminate inflorescence is an umbellate cluster of spikes 3-5.5 cm long, consisting of many individual tubular calyces with paired stamens, while its pistillate spikes are yellowish and 2-5.5 cm long, thick, and succulent when in fruit (PIER, 2009).

Notes

Cecropia peltata L was distinguished from C. schreberiana Mig. in 1988. Whereas Cecropia peltata occurs in Mexico and Central America, C. schreberiana occurs in the Antilles and northern South America (Howard, 1988; ISTF, 1997 in Brokaw, 1998; Csuhres, 2008). However, ITIS does not distinguish between the species and, in fact, states Cecropia schreberiana as the valid name for the species and indicates C. peltata as a synonym for C. schreberiana.



FULL ACCOUNT FOR: Cecropia peltata

Lifecycle Stages

Seeds of *Cecropia peltata* require full sunlight for successful germination and with those conditions may be as high as 80-90%. Seedling leaves are pubescent on both sides, lanceolate, unlobed, and finely toothed. Seedlings are also very light demanding and seedling mortality in natural conditions is typically very high. It has been found that 99% of seedlings in forest openings die in the first year. *C. peltata* grows rapidly reaching 10-15 cm in height in 10 weeks and up to about 2 m in the first year. Reproductive maturity is reached by pistillate trees in 3-4 years and by staminate trees in 4-5 years. Maturation is dependant on allocation of resources for rapid initial height growth and factors such as the height of and proximity to surrounding vegetation with trees in open environments maturing faster than those in forest gaps. *C. peltata* usually reaches canopy height in about 10 years and its estimated life span is 30 years (Silander & Lugo, undated).

Uses

Cecropia peltata is popularly cultivated as an ornamental species (Bodkin 1990 in Csurhes, 2008).

Habitat Description

Cecropia peltata typically inhabits forest gaps and disturbed sites (PIER, 2009), such as, along roadsides, agricultural sites, lava flows, and urban locations (Binggeli, 1999). It is a fast growing, high light demanding, pioneer species that colonizes tree fall gaps in ts native range and is capable of establishing dense stands (PIER, 2009). It is known from altitudes of 50-2700 m (Hurtado & Alson, 1995). *C. peltata* requires much rainfall and may be found in environments with 990 mm to over 3,810 mm of annual percipitation. It grows in alluvial, colluvial, and residual soils neutral to acidic in nature. Soil texture may range from heavy clay to sandy, but a clay-loam soil is optimal. *C. peltata* is also generally found in warm climates ranging from montane to tropical with mean annual temperatures of 12-24°C (Silander & Lugo, undated).

Reproduction

Cecropia peltata is dioecious and becomes sexually mature in 3 to 5 years. Its tiny flowers are clustered on 5 to 10 cm long spikes and are wind-pollinated. On female spikes the minute one-seeded fruits form large fruit clusters which appear to take around a month to mature. A spike contains around 800 viable seeds which are about 1.9 mm long and weigh 1.6 mg. Bats and birds eat large quantities of the succulent fruits and are the main seed disperser. In Costa Rica a similar amount of fruits are consumed during the day, mainly by monkeys, and at night by bats and arboreal mammals. A large and persistent seedbank is formed in the forest soil (Bingelli, 1999). In some locations flowering and fruiting occur year-round and in others it it seasonal with a peak in either the wet or the dry season depending on location (Silander & Lugo, undated; Bingelli, 1999). C. peltata is highly productive and seed production is estimated to be as high as 1 million seeds per year (Silander & Lugo, undated)

General Impacts

Cecropia peltata forms dense stands that may compete with or displace native pioneer species and reduce species richness (Bingelli, 1999; Dumont et al, 1990). Evidence suggests it competes with and may displace tropical African pioneer species *Musanga cecropioides* (Bingelli, 1999).



FULL ACCOUNT FOR: Cecropia peltata

Management Info

<u>Preventative measures</u>: A Risk Assessment of *Cecropia peltata* for Hawai'i and other Pacific islands was prepared by Dr. Curtis Daehler (UH Botany) with funding from the Kaulunani Urban Forestry Program and US Forest Service. The alien plant screening system is derived from Pheloung *et al.* (1999) with minor modifications for use in Pacific islands (Daehler *et al.* 2004). The result is a score of 9 and a recommendation of: \"Likely to cause significant ecological or economic harm in Hawai'i and on other Pacific Islands as determined by a high WRA score, which is based on published sources describing species biology and behavior in Hawai'i and/or other parts of the world.\"

\nPhysical: Hand pulling or digging out seedlings and young trees is recommended (PIER, 2009). \nChemical: Larger trees should be cut and their stumps should be treated with herbicide (PIER, 2009). \nBiological control: C. peltata has been found to be attacked by Historis spp. and various moth species and is sometimes extensively defoliated (Bingelli et al, 1998). \n

Pathway

Principal source: Bingelli, Pierre, 1999. *Cecropia peltata* L. (Cecropiaceae). Pacific Ecosystems at Risk (PIER), 2009. *Cecropia peltata* Regel, Cecropiaceae

Csurhes, Steve, 2008. *Cecropia*, *Cecropia* spp. Pest Plant Risk Assessment. Biosecurity Queensland Department of Primary Industries and Fisheries, Queensland

Silander, Susan R. and Ariel E. Lugo, undated. Cecropia peltata L. Yagrumo Hembra, Trumpet-Tree.

Compiler: National Biological Information Infrastructure (NBII) & IUCN/SSC Invasive Species Specialist Group (ISSG)

Review:

Pubblication date: 2011-02-23

ALIEN RANGE

[1] ANGOLA[1] AUSTRALIA[1] CAMEROON[1] COTE D'IVOIRE[7] FRENCH POLYNESIA[1] MADAGASCAR[1] MALAYSIA[1] NEW CALEDONIA

BIBLIOGRAPHY

25 references found for Cecropia peltata

Managment information

<u>Australian Weed Committee</u>, 2010. Weed Identification: Australia > > Trumpet Tree

Summary: Available from: http://www.weeds.org.au/cgi-bin/weedident.cgi?tpl=plant.tpl&state=&s=&ibra=all&card=T38 [Accessed March 25 2010]

Binggeli, Pierre; John B. Hall and John R. Healy, 1998. An Overview of Invasive Woody Plants in the Tropics

Summary: Available from: http://www.bangor.ac.uk/~afs101/iwpt/welcome.shtml [Accessed March 25 2010]

Csurhes, Steve, 2008. Cecropia, Cecropia spp. Pest Plant Risk Assessment. Biosecurity Queensland Department of Primary Industries and Fisheries, Queensland

Daehler, C.C; Denslow, J.S; Ansari, S and Huang-Chi, K., 2004. A Risk-Assessment System for Screening Out Invasive Pest Plants from Hawaii and Other Pacific Islands. Conservation Biology Volume 18 Issue 2 Page 360.

Summary: A study on the use of a screening system to assess proposed plant introductions to Hawaii or other Pacific Islands and to identify high-risk species used in horticulture and forestry which would greatly reduce future pest-plant problems and allow entry of most nonpests. Pacific Ecosystems at Risk (PIER), 2009. Cecropia peltata Regel, Cecropiaceae

Summary: Available from: http://www.hear.org/pier/species/cecropia_peltata.htm [Accessed March 25 2010]

Pacific Ecosystems at Risk (PIER), 2009. Risk Assessment Cecropia peltata Regel, Cecropiaceae

Summary: Available from: http://www.hear.org/pier/wra/pacific/cecropia peltata htmlwra.htm [Accessed March 25 2010]

General information



FULL ACCOUNT FOR: Cecropia peltata

Bingelli, Pierre, 1999. Cecropia peltata L. (Cecropiaceae)

Summary: Available from: http://www.bangor.ac.uk/~afs101/iwpt/web-sp3.htm [Accessed March 25 2010]

CABI, Forestry Compendium, 2006. Common names Cecropia peltata

Summary: Available from: http://www.cabicompendium.org/NamesLists/FC/Full/CECRPE.htm [Accessed March 25 2010]

Dumont, J.F., Lamotte, S. and Kahn, F., 1990. Wetland and upland forest ecosystems in Peruvian Amazonia: Plant species diversity in the light of some geological and botanical evidence. For. Ecol. Manage., 33/34: 125-139.

Summary: Available from: http://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_5/b_fdi_30-30/31531.pdf [Accessed March 25 2010]

Fleming, Theodore H. and Charles F. Williams, 1990. Journal of Tropical Ecology (1990), 6: 163-178 Cambridge University Press Florence J. Chevillotte H. Ollier C.& Meyer J.-Y., 2007. Cecropia peltata. Base de données botaniques Nadeaud de l Herbier de la Polynésie franéaise (PAP).

Summary: Base de donn es sur le flore de Polyn sie Fran aise.

Available from: http://www.herbier-tahiti.pf/Selection_Taxonomie.php?id_tax=9466 [Accessed March 2008]

Féveri, Sarita B. & Heraldo L. Vasconcelos, 2004. The Azteca Cecropia Association: Are Ants Always Necessary for Their Host Plants? Biotropica Dec 2004: Vol. 36, Issue 4, pg(s) 641-646 doi: 10.1646/1606

Howard, R. A. 1988. Flora of the Lesser Antilles, volume 4. Arnold Arboretum, Jamaica Plains, MA.

Hurtado, In s and Julio Alson, 1995. Airborne dispersal of *Cecropia peltata* L. (Moraceae) pollen, a neotropical species, in the vicinity of Caracas, Venezuela. Aerobiologia 11 (1995) 101-103

IABIN, 2008. Guarumo Species and Specimens Thematic Network

Summary: Available from: http://ara.inbio.ac.cr/SSTN-IABIN/species/157260 [Accessed March 25 2010]

Integrated Taxonomic Information System (ITIS), 2010. Cecropia schreberiana Mig.

Summary: Available from: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=565052 [Accessed March 25 2010]

James Cook University, 2010. Cecropia peltata

Summary: Available from: http://cms.jcu.edu.au/discovernature/weeds/JCUPRD 043283 [Accessed March 25 2010]

MacKee, H.S. 1994. Catalogue des plantes introduites et cultivêes en Nouvelle-Calêdonie, 2nd edn. MNHN, Paris.

Summary: Cet ouvrage liste 1412 taxons (espêces, sous espêces et variêtês) introduits en Nouvelle-Calêdonie. L auteur prêcise dans la majoritê des cas si l espêce est cultivêe ou naturalisêe.

McKey, Doyle, 1988. *Cecropia peltata*, an Introduced Neotropical Pioneer Tree, is Replacing *Musanga cecropioides* in Southwestern Cameroon. Biotropica, Vol. 20, No. 3 (Sep., 1988), pp. 262-264

Meyer, Jean-Yves and Jacques Florence, 1996. Tahiti s Native Flora Endangered by the invasion of *Miconia calvescens* DC Melastomataceae. Journal of Biogeography (1996) 23, 775-781

Summary: Available from: http://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_7/b_fdi_51-52/010014779.pdf [Accessed March 25 2010]

Meyer, Jean-Yves & Loope, Lloyd & Sheppard, A. & Munzinger, Jérôme & Jaffré, Tanguy. (2006). Les plantes envahissantes et potentiellement envahissantes dans l'archipel néo-calédonien : première évaluation et recommandantions de gestion.

Meyer, J.-Y. 2004. Threat of invasive alien plants to native flora and forest vegetation of eastern Polynesia. Pacific Science, 58, 357-375

Summary: Dans cet article, la menace croissante des plantes exotiques envahissantes est discutê et les espêces les plus envahissantes sont décrites. Des hypothèses sur l invasibilitê des éles sont présentées à la lumière des observations et des données récoltées. Silander, S. R. and Lugo, A. E. Cecropia peltata L. (Yagrumo Hembra, Trumpet-Tree). Silvics of North American, USDA Agriculture Handbook

Summary: Fact sheet.

Silander, Susan R. and Ariel E. Lugo, undated. Cecropia peltata L. Yagrumo Hembra, Trumpet-Tree

Summary: Available from: http://www.na.fs.fed.us/pubs/silvics_manual/Volume_2/cecropia/peltata.htm [Accessed March 25 2010] USDA, NRCS. 2010. *Cecropia schreberiana* Miq. Pumpwood. The PLANTS Database (http://plants.usda.gov, 26 March 2010). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

Summary: Available from: http://plants.usda.gov/java/profile?symbol=CESC9 [Accessed March 25 2010]