Batrachochytrium dendrobatidis 简体中文 正體中文

System: Undefined

Kingdom	Phylum	Class	Order	Family
Fungi	Chytridiomycota	Chytridiomycetes	Chytridiales	

Common name chytrid frog fungi (English), Chytrid-Pilz (German), chytridiomycosis (English), frog chytrid fungus (English)

Synonym

Similar species

Summary Batrachochytrium dendrobatidis is a non-hyphal parasitic chytrid fungus that has been associated with population declines in endemic amphibian species in upland montane rain forests in Australia and Panama. It causes cutaneous mycosis (fungal infection of the skin), or more specifically chytridiomycosis, in wild and captive amphibians. First described in 1998, the fungus is the only chytrid known to parasitise vertebrates. *B. dendrobatidis* can remain viable in the environment (especially aquatic environments) for weeks on its own, and may persist in latent infections.

view this species on IUCN Red List

Species Description
Fungal Morphology: *Batrachochytrium dendrobatidis* is a zoosporic chytrid fungus that causes chytridiomycosis (a fungal infection of the skin) in amphibians and grows solely within keratinised cells. Diagnosis is by identification of characteristic intracellular flask-shaped sporangia (spore containing bodies) and septate thalli. The fungus grows in the superficial keratinised layers of the epidermis (known as the stratum corneum and stratum granulosum). The normal thickness of the stratum corneum is between 2µm to 5µm, but a heavy infection by the chytrid parasite may cause it to thicken to up to 60 µm. The fungus also infects the mouthparts of tadpoles (which are keratinised) but does not infect the epidermis of tadpoles (which lacks keratin).
The fungus produces inoperculate, smooth-walled zoosporangia (zoospore containing bodies), which are spherical to subspherical in shape. Each zoosporangium (10µm to 40µm in diameter) produces a single discharge tube, which penetrates (and protrudes out of) the skin. Eventually the plug that blocks the release of immature zoospores is shed and the mature zoospores are released. The zoospores (0.7µm to 6µm in diameter) are elongate to ovoid in shape. Each possesses a single posterior flagellum, rendering it motile in water (Mazzoni *et al.* 2003; Daszak *et al.* 1999; Berger, *et al.* 1998; Berger *et al.* 1998, Berger, Speare and Hyatt, 2000, in Daszak *et al.* 1999; Speare *et al.* 2001; Weldon *et al.* 2003).\r\n
To view a scanning electron micrograph of infected skin of a wild frog (*Litoria lesueuri*) please see: \r\n *Daszak et al. 1999. Emerging Infectious Diseases and Amphibian Population Declines*.\r\n
To view histological sections of infected skin of *Bufo haematiticus* and *Atelopus varius* (showing the sporangia and discharge tubes of the fungus) please see: *Daszak et al. 1999. Emerging Infectious Diseases and Amphibian Population Declines*.\r\n
To view a histological section of severely infected skin of a wild frog (*Litoria caerulea*) please see:\r\n *Berger et al. 1998. Chytridiomycosis causes amphibian mortality* .
\r\nClick here to see information about Symptoms of the disease caused by *Batrachochytrium dendrobatidis*.
Pathogenesis of chytridiomycosis: Authors of a recent study, Voyles *et al*. (2009) have found that *B. dendrobatidis*, causes such severe electrolyte imbalances that the frog's heart stops. The skin of amphibians maintain proper osmotic balance inside the animal and regulate respiration. The authors found that the skin of infected frogs was less adept at transporting sodium and chloride ions. Sodium and potassium concentrations in the blood of infected frogs dropped, more so as the infection intensified and the animals'hearts began to beat irregularly and ultimately stopped.

Notes
Salamanders can act as host reservoirs of chytrid infection in frogs, and vice versa (Davidson *et al.* 2003).

Lifecycle Stages
Batrachochytrium dendrobatidis has two life stages: a spherical reproductive sessile zoosporangium and a motile zoospore. The motile zoospore directs itself and attaches to the keratinised outer layers of its host. It then absorbs its tail and buries itself below the surface of the skin. It matures into a zoosporangia with rhizoids within about four days and produces and releases up to 300 zoospores into the external environment (*via* a discharge tube). The cycle is initiated again once a suitable substrate (in the same or a different host) is found. The presence of the fungus in the keratinised mouthparts of frog tadpoles (without actually killing them) supports the role of larvae as reservoirs for the pathogen. (The larvae of amphibian species may survive for as long as 3 years before metamorphosing.) Syntopic salamanders and frogs may also act as reciprocal pathogen reservoirs for chytrid infections. It has been suggested that *B. dendrobatidis* may not be an obligate amphibian parasite, possibly living in other non-amphibian hosts or even sapropytically (off dead tissue) (Michigan Frog Survey, 2003; Speare *et al.* 2001; Daszak *et al.* 1999; Davidson *et al.* 2003).\n
As of yet, no resting structures (either asexual or sexual) have been identified for *B. dendrobatidis*. The fact that sexual reproduction in chytrid fungi has been associated with the production of resistant, thick-walled resting spores has lead to the hypothesis that the production of airborne spores explains the widespread distribution of *B. dendrobatidis* in relatively pristine areas. However recent research has found evidence that shows that the population structure of *B. dendrobatidis* is largely clonal, supporting the hypothesis that the fungus lacks a sexual stage (as is the case for many chytrid fungi). This suggests that dispersal by human (or perhaps other long distance travellers, such as birds), rather than natural causes, are more likely to be the cause of the pathogen's entry into pristine areas (Morehouse *et al.* 2003; Berger *et al.* 1999, Daszak *et al.* 1999, in Morehouse *et al.* 2003).\"

Habitat Description
Chytridiomycosis has now been reported from 38 amphibian species in 12 families, including ranid and hylid frogs, bufonid toads, and plethodontid salamanders. Although chytridiomycosis is found in a range of species and habitats (including African frogs in lowland regions in Africa) it has caused population declines of amphibians species confined to montane rain forests (Weldon *et al.* 2004; Daszak *et al.* 1999). The fungus prefers lower temperatures which may explain the high precedence of the fungus in high elevations in the tropics. In culture conditions optimum growth occurred at 23°C, with slower growth occuring at 28°C and (reversible) cessation of growth occuring at 29°C (Longcore, Pessier, Nichols, 1999, in Daszak *et al.* 1999).

Reproduction
Batrachochytrium dendrobatidis is diploid and primarily reproduces asexually (and clonally) by producing aquatic uniflagellated zoospores in a zoosporangium (Johnson and Speare, 2003).

Nutrition
Its occurrence solely in keratinised tissues suggests that it uses amphibian keratin as a nutrient. *Batrachochytrium dendrobatidis* will grow for at least one generation on cleaned epidermal keratin or on amphibians that have died of the infection. The fungus may also be cultured *in vitro* on tryptone agar without the addition of keratin or its derivatives (Daszak *et al.* 1999; Longcore, Pessier and Nichols, 1999, Pessier *et al.* 1999, in Daszak *et al.* 1999).

General Impacts
Batrachochytrium dendrobatidis has been found to affect at least 93 amphibian species from the orders Anura (frogs and toads) and Caudata (salamanders) in all the continents except Asia. It is thought to be one of the main causes of the global decline in frog populations since the 1960s, and the dramatic population crashes from the 1970s onwards (Parris and Beaudoin, 2004). The chytrid fungus kills frogs within 10 to 18 days (Michigan Frog Survey, 2003), although it is not known how. It may be physical, affecting respiration by altering the frog's skin, or the fungus may give off a toxin (Michigan Frog Survey, 2003). Tadpoles are not affected, although the fungus may infect the keratinised mouthparts (Berger *et al.* 1999).
For a summary on the impacts of *B. dendrobatidis* please follow this link *impacts*.
Key findings of the *The Global Amphibian Assessment* has revealed that one-third (32%) of the world's amphibian species are threatened, representing 1,896 species. Threats include viral diseases, habitat loss, drought, pollution, and hunting for food. The biggest single threat appears to be *B. dendrobatidis*.
A *search* on the database using \"diseases\" as a keyword in \"all\" habitat types, biogeographic realm and countries results in a list of 547 species impacted by diseases (IUCN, Conservation International, and NatureServe. 2006).

Management Info
Preventative measures: Knowledge of the infectiveness and spread of *Batrachochytrium dendrobatidis* is relevant to all control strategies, particularly in the development of preventative measures. The infective unit of the fungus is the zoospore. Infection by the fungus (and thus spread of the disease) requires water because the zoospore does not tolerate dehydration. *B. dendrobatidis* remains viable for up to 3 weeks in tap water, up to 4 weeks in deionised water and even longer in lake water. Infection by an extremely small inoculum (100 zoospores) is sufficient to cause a fatal infection (Berger *et al.* in Speare *et al.* 2001; Johnson and Speare, 2003; Berger, Speare and Hyatt, 2000, in Daszak *et al.* 1999).
Please see *main preventative management strategies* for a summary under the following headings: improving diagnostics and knowledge of epidemiology, developing trade and quarantine regulations, raising awareness and control options.
The Amphibian Conservation Action Plan (ACAP) is designed to provide guidance for implementing amphibian conservation and research initiatives at all scales from global down to local. Chapter 4 outlines action steps relating to the detection and control of chytridiomycosis.

Principal source: Berger *et al.* 1999. Chytrid fungi and amphibian declines: Overview, Implications and Future Directions.
Berger *et al.* 1998. Chytridiomycosis Causes Amphibian Mortality Associated With Population Declines in the Rain Forests of Australia and Central America.
Daszak *et al.* 1999. Emerging Infectious Diseases and Amphibian Population Declines

Compiler: National Biological Information Infrastructure (NBII) & IUCN/SSC Invasive Species Specialist Group (ISSG) with support from the Terrestrial and Freshwater Biodiversity Information System (TFBIS) Programme (Copyright statement)

Review: Matthew J. Parris Assistant Professor, Department of Biology University of Memphis USA

Pubblication date: 2006-08-14

ALIEN RANGE

[85] AUSTRALIA
[4] COSTA RICA
[2] GERMANY
[1] ITALY
[3] MEXICO
[2] CANADA
[4] ECUADOR
[1] GHANA
[2] KENYA
[5] NEW ZEALAND

[3] PANAMA
[1] SPAIN
[14] UNITED STATES
[1] VENEZUELA
[9] SOUTH AFRICA
[1] SWAZILAND
[2] URUGUAY
[1] WEST AFRICA

Red List assessed species 512: EX = 8; CR = 196; EN = 126; VU = 63; NT = 29; DD = 36; LC = 54;

Adelotus brevis **NT**
Agalychnis moreletii **CR**
Alytes cisternasii **NT**
Aplastodiscus callipygius **LC**
Aromobates alboguttatus **EN**
Aromobates nocturnus **CR**
Atelopus angelito **CR**
Atelopus arthuri **CR**
Atelopus bomolochos **CR**
Atelopus carauta **CR**
Atelopus carrikeri **CR**
Atelopus chiriquiensis **CR**
Atelopus chrysocorallus **CR**
Atelopus cruciger **CR**
Atelopus ebenoides **CR**
Atelopus epikeisthos **CR**
Atelopus eusebianus **CR**
Atelopus famelicus **CR**
Atelopus flavescens **VU**
Atelopus galactogaster **CR**
Atelopus guanujo **CR**
Atelopus halihelos **CR**
Atelopus laetissimus **CR**
Atelopus longibrachius **EN**
Atelopus lozanoi **CR**
Atelopus mandingues **CR**
Atelopus minutulus **CR**
Atelopus monohernandezii **CR**
Atelopus muisca **CR**
Atelopus nanay **CR**
Atelopus nicefori **CR**
Atelopus oxapampae **EN**
Atelopus pachydermus **CR**
Atelopus patazensis **CR**
Atelopus peruensis **CR**
Atelopus petriruizi **CR**
Atelopus pinangoi **CR**
Atelopus pulcher **CR**
Atelopus reticulatus **CR**
Atelopus seminiferus **CR**
Atelopus sernai **CR**
Atelopus siranus **DD**
Atelopus sorianoi **CR**
Atelopus spurrelli **VU**
Atelopus tamaense **CR**
Atelopus varius **CR**
Atelopus zeteki **CR**
Bokermannohyla claresignata **DD**
Agalychnis annae **EN**
Allobates olfersioides **VU**
Anaxyrus canorus **EN**
Aplastodiscus flumineus **DD**
Aromobates leopardalis **CR**
Atelopus andinus **CR**
Atelopus arsyecue **CR**
Atelopus balios **CR**
Atelopus boulengeri **CR**
Atelopus carbonerensis **CR**
Atelopus certus **EN**
Atelopus chocoensis **CR**
Atelopus coynei **CR**
Atelopus dimorphus **EN**
Atelopus elegans **CR**
Atelopus erythropus **CR**
Atelopus exiguus **CR**
Atelopus farci **CR**
Atelopus franciscus **VU**
Atelopus glyphus **CR**
Atelopus guitarraensis **CR**
Atelopus ignescens **EX**
Atelopus limosus **EN**
Atelopus longirostris **EX**
Atelopus lynchi **CR**
Atelopus mindoensis **CR**
Atelopus mittermeieri **EN**
Atelopus mucubajiensis **CR**
Atelopus nahumae **CR**
Atelopus nepiozomus **CR**
Atelopus onorei **CR**
Atelopus oxyrhynchus **CR**
Atelopus palmatus **DD**
Atelopus pedimarmoratus **CR**
Atelopus petersi **CR**
Atelopus pictiventris **CR**
Atelopus planispina **CR**
Atelopus quimbaya **CR**
Atelopus sanjosei **DD**
Atelopus senex **CR**
Atelopus simulatus **CR**
Atelopus sonsonensis **CR**
Atelopus spumarius **VU**
Atelopus subornatus **CR**
Atelopus tricolor **VU**
Atelopus walkeri **CR**
Bokermannohyla circumdata **LC**
Bokermannohyla hylax **LC**

Bolitoglossa conanti **EN**
Bolitoglossa dofleini **NT**
Bolitoglossa pesrubra **VU**
Bolitoglossa sooyorum **EN**
Bombina pachypus **EN**
Bromeliohyla dendroscarta **CR**
Centrolene audax **EN**
Centrolene buckleyi **VU**
Centrolene gemmatum **CR**
Centrolene lynchi **EN**
Centrolene peristictum **VU**
Centrolene scirtetes **DD**
Charadrahyla nephila **VU**
Chiropterotriton cracens **EN**
Craugastor anciano **CR**
Craugastor angelicus **CR**
Craugastor berkenbuschii **NT**
Craugastor catalinae **CR**
Craugastor chrysozetetes **EX**
Craugastor daryi **EN**
Craugastor emleni **CR**
Craugastor escoces **EX**
Craugastor fleischmanni **CR**
Craugastor guerreroensis **CR**
Craugastor laevissimus **EN**
Craugastor lineatus **CR**
Craugastor merendonensis **CR**
Craugastor milesi **CR**
Craugastor olanchano **CR**
Craugastor pechorum **EN**
Craugastor podiciferus **NT**
Craugastor punctariolus **EN**
Craugastor rhyacobatrachus **EN**
Craugastor rugulosus **LC**
Craugastor sabrinus **EN**
Craugastor sandersoni **EN**
Craugastor tabasarae **CR**
Craugastor trachydermus **CR**
Crinia pseudinsignifera **LC**
Crossodactylus gaudichaudii **LC**
Cycloramphus ohausi **DD**
Duellmanohyla chamulae **EN**
Duellmanohyla lythrodes **EN**
Duellmanohyla schmidtorum **VU**
Duellmanohyla uranochroa **CR**
Ecnomiohyla rabborum **CR**
Eleutherodactylus cooki **VU**
Eleutherodactylus hedricki **EN**
Eleutherodactylus karlschmidti **CR**
Eleutherodactylus orcutti **CR**
Eleutherodactylus portoricensis **EN**
Eleutherodactylus ruthae **EN**
Eleutherodactylus semipalmatus **CR**
Eleutherodactylus turquinensis **CR**
Bolitoglossa copia **DD**
Bolitoglossa magnifica **EN**
Bolitoglossa sombra **VU**
Bolitoglossa subpalmata **EN**
Bromeliohyla bromeliacia **EN**
Bufo bufo **LC**
Centrolene ballux **CR**
Centrolene geckoideum **VU**
Centrolene heloderma **CR**
Centrolene medemi **DD**
Centrolene pipilatum **EN**
Charadrahyla altipotens **CR**
Charadrahyla trux **CR**
Chiropterotriton multidentatus **EN**
Craugastor andi **CR**
Craugastor azueroensis **EN**
Craugastor brocchi **VU**
Craugastor charadra **EN**
Craugastor cruzi **CR**
Craugastor emcelae **CR**
Craugastor epochthidius **CR**
Craugastor fecundus **CR**
Craugastor greggi **CR**
Craugastor inachus **EN**
Craugastor laticeps **NT**
Craugastor melanostictus **LC**
Craugastor mexicanus **LC**
Craugastor obesus **EN**
Craugastor omoaensis **CR**
Craugastor phasma **DD**
Craugastor polymniae **CR**
Craugastor ranoides **CR**
Craugastor rostralis **NT**
Craugastor rupinius **LC**
Craugastor saltuarius **CR**
Craugastor stadelmani **CR**
Craugastor taurus **CR**
Crinia georgiana **LC**
Crossodactylus dispar **DD**
Cycloramphus boraceiensis **LC**
Cycloramphus semipalmatus **NT**
Duellmanohyla ignicolor **EN**
Duellmanohyla salvavida **CR**
Duellmanohyla soralia **CR**
Ecnomiohyla echinata **CR**
Eleutherodactylus barlagnei **EN**
Eleutherodactylus gryllus **EN**
Eleutherodactylus jasperi **CR**
Eleutherodactylus longipes **VU**
Eleutherodactylus patriciae **EN**
Eleutherodactylus richmondi **CR**
Eleutherodactylus schmidti **CR**
Eleutherodactylus symingtoni **CR**
Eleutherodactylus unicolor **VU**

Eleutherodactylus wightmanae **EN**
Euproctus platycephalus **EN**
Exerodonta melanomma **VU**
Gastrotheca cornuta **EN**
Gastrotheca guentheri **VU**
Gastrotheca orophylax **EN**
Gastrotheca piperata **LC**
Gastrotheca pseustes **EN**
Gastrotheca splendens **EN**
Heleioporus australiacus **VU**
Hyalinobatrachium fleischmanni **LC**
Hyla bocourti **CR**
Hylodes dactylocinus **DD**
Hylodes meridionalis **LC**
Hylodes phyllodes **LC**
Hyloscirtus armatus **LC**
Hyloscirtus colymba **CR**
Hyloscirtus pantostictus **EN**
Hyloscirtus ptychodactylus **CR**
Hyloscirtus torrenticola **VU**
Hyloxalus bocagei **LC**
Hyloxalus chocoensis **DD**
Hyloxalus elachyhistus **EN**
Hyloxalus pulchellus **VU**
Hypodactylus dolops **VU**
Incilius cycladen **VU**
Incilius holdridgei **EX**
Incilius periglenes **EX**
Incilius porteri **DD**
Incilius tutelarius **EN**
Isthmohyla calypsa **CR**
Isthmohyla graceae **CR**
Isthmohyla tica **CR**
Leiopelma archeyi **CR**
Leiopelma hochstetteri **VU**
Leptodactylus fallax **CR**
Lithobates chiricahuensis **VU**
Lithobates sierramadrensis **VU**
Lithobates tarahumarae **VU**
Lithobates vibicarius **CR**
Lithobates yavapaiensis **LC**
Litoria aurea **VU**
Litoria caerulea **LC**
Litoria chloris **LC**
Litoria ewingii **LC**
Litoria lesueurii **LC**
Litoria moorei **LC**
Litoria nannotis **EN**
Litoria nyakalensis **CR**
Litoria phyllochroa **LC**
Litoria raniformis **EN**
Litoria spenceri **CR**
Mannophryne caquetio **CR**
Mannophryne herminae **NT**
Epipedobates tricolor **EN**
Exerodonta juanitae **VU**
Exerodonta pinorum **VU**
Gastrotheca dendronastes **VU**
Gastrotheca litonedis **EN**
Gastrotheca ovifera **EN**
Gastrotheca plumbea **VU**
Gastrotheca riobambae **EN**
Geocrinia rosea **LC**
Heleioporus eyrei **LC**
Hyalinobatrachium guairarepanense **EN**
Hylarana chalconota **LC**
Hylodes magalhaesi **DD**
Hylodes perplicatus **LC**
Hylomantis lemur **CR**
Hyloscirtus bogotensis **NT**
Hyloscirtus lindae **VU**
Hyloscirtus platydactylus **VU**
Hyloscirtus staufferorum **EN**
Hyloxalus anthracinus **CR**
Hyloxalus breviquartus **DD**
Hyloxalus delatorreae **CR**
Hyloxalus lehmanni **NT**
Hyloxalus vertebralis **CR**
Hypsiboas cymbalum **CR**
Incilius fastidiosus **CR**
Incilius melanochlorus **LC**
Incilius peripatetes **CR**
Incilius tacanensis **EN**
Isthmohyla angustilineata **CR**
Isthmohyla debilis **CR**
Isthmohyla pictipes **EN**
Isthmohyla xanthosticta **DD**
Leiopelma hamiltoni **EN**
Leptobrachium hasseltii **LC**
Limnodynastes dumerilii **LC**
Lithobates omiltemanus **CR**
Lithobates subaquavocalis **CR**
Lithobates taylori **LC**
Lithobates warszewitschii **LC**
Litoria adelaidensis **LC**
Litoria booroolongensis **CR**
Litoria castanea **CR**
Litoria dayi **EN**
Litoria genimaculata **LC**
Litoria lorica **CR**
Litoria myola **CR**
Litoria nudidigita **LC**
Litoria pearsoniana **NT**
Litoria piperata **CR**
Litoria rheocola **EN**
Litoria verreauxii **LC**
Mannophryne cordilleriana **CR**
Mannophryne lamarcai **CR**

Mannophryne neblina **CR**
Mannophryne olmonae **CR**
Megaelosia massarti **DD**
Mesotriton alpestris **LC**
Mixophyes fasciolatus **LC**
Nymphargus griffithsi **VU**
Oedipina **EN**
Osteopilus pulchrilineatus **EN**
Paratelmatobius lutzii **DD**
Pelobates fuscus **LC**
Phrynomedusa appendiculata **NT**
Phyllomedusa ecuatoriana **EN**
Physalaemus moreirae **DD**
Plectrohyla ameibothalame **DD**
Plectrohyla avia **CR**
Plectrohyla calthula **CR**
Plectrohyla celata **CR**
Plectrohyla charadricola **EN**
Plectrohyla chrysopleura **CR**
Plectrohyla cyanomma **CR**
Plectrohyla dasypus **CR**
Plectrohyla exquisita **CR**
Plectrohyla guatemalensis **CR**
Plectrohyla hazelae **CR**
Plectrohyla lacertosa **EN**
Plectrohyla mykter **EN**
Plectrohyla pentheter **EN**
Plectrohyla psiloderma **EN**
Plectrohyla quecchi **CR**
Plectrohyla sabrina **CR**
Plectrohyla siopela **CR**
Plectrohyla teuchestes **CR**
Pleurodema marmoratum **LC**
Pristimantis anotis **DD**
Pristimantis calcarulatus **VU**
Pristimantis caryophyllaceus **NT**
Pristimantis cremnobates **EN**
Pristimantis crucifer **VU**
Pristimantis diogenes **VU**
Pristimantis fallax **EN**
Pristimantis ginesi **EN**
Pristimantis ignicolor **EN**
Pristimantis jorgevelosai **EN**
Pristimantis lancinii **EN**
Pristimantis lymani **LC**
Pristimantis nigrogriseus **VU**
Pristimantis prolatus **EN**
Pristimantis ruedai **VU**
Pristimantis sanguineus **NT**
Pristimantis scoloblepharus **EN**
Pristimantis signifer **VU**
Pristimantis tamsitti **NT**
Pristimantis urichi **EN**
Pristimantis vicarius **NT**
Mannophryne oblitterata **DD**
Mannophryne riveroi **EN**
Megastomatohyla pellita **CR**
Mixophyes balbus **VU**
Mixophyes fleayi **EN**
Nymphargus megacheirus **EN**
Oophaga arborea **EN**
Osteopilus vastus **EN**
Paratelmatobius mantiqueira **DD**
Philoria frosti **CR**
Phyllobates bicolor **NT**
Physalaemus barrioi **DD**
Plectrohyla acanthodes **CR**
Plectrohyla arborescandens **EN**
Plectrohyla bistincta **LC**
Plectrohyla calvicollina **CR**
Plectrohyla cembra **CR**
Plectrohyla chryses **CR**
Plectrohyla crassa **CR**
Plectrohyla cyclada **EN**
Plectrohyla ephemera **CR**
Plectrohyla glandulosa **EN**
Plectrohyla hartwegi **CR**
Plectrohyla ixil **CR**
Plectrohyla matudai **VU**
Plectrohyla pachyderma **CR**
Plectrohyla pokomchi **CR**
Plectrohyla pycnochila **CR**
Plectrohyla robertsorum **EN**
Plectrohyla sagorum **EN**
Plectrohyla tecunumani **CR**
Plectrohyla thorectes **CR**
Pristimantis albericoi **CR**
Pristimantis bicolor **VU**
Pristimantis caprifer **LC**
Pristimantis chalceus **LC**
Pristimantis crenunguis **EN**
Pristimantis diaphonus **VU**
Pristimantis duellmani **VU**
Pristimantis fetosus **EN**
Pristimantis gracilis **VU**
Pristimantis incanus **EN**
Pristimantis labiosus **LC**
Pristimantis lichenoides **CR**
Pristimantis molybrignus **NT**
Pristimantis penelopus **VU**
Pristimantis quinquagesimus **VU**
Pristimantis sanctaemartae **NT**
Pristimantis savagei **NT**
Pristimantis scolodiscus **EN**
Pristimantis sulculus **EN**
Pristimantis uranobates **LC**
Pristimantis verecundus **VU**
Pristimantis zophus **EN**

Prostherapis dunni **CR**
Pseudoeurycea unguidentis **CR**
Pseudophryne pengilleyi **EN**
Ptychohyla acrochorda **DD**
Ptychohyla erythromma **EN**
Ptychohyla legleri **EN**
Ptychohyla macrotympanum **CR**
Ptychohyla salvadorensis **EN**
Ptychohyla spinipollex **EN**
Rana muscosa **EN**
Ranitomeya abdita **CR**
Rhaebo haematiticus **LC**
Rhinella amabilis **CR**
Rhinoderma darwinii **VU**
Scinax albicans **LC**
Silverstoneia nubicola **NT**
Strabomantis cheiroplethus **VU**
Strabomantis zygodactylus **LC**
Taudactylus diurnus **EX**
Taudactylus liemi **NT**
Taudactylus rheophilus **CR**
Telmatobius atacamensis **CR**
Telmatobius bolivianus **NT**
Telmatobius brevirostris **EN**
Telmatobius ceiorum **EN**
Telmatobius colanensis **EN**
Telmatobius culeus **CR**
Telmatobius degener **EN**
Telmatobius gigas **CR**
Telmatobius hockingi **VU**
Telmatobius ignavus **EN**
Telmatobius jelskii **NT**
Telmatobius latirostris **EN**
Telmatobius mayoloi **EN**
Telmatobius niger **CR**
Telmatobius peruvianus **VU**
Telmatobius pinguiculus **DD**
Telmatobius platycephalus **EN**
Telmatobius scrocchii **EN**
Telmatobius simonsi **NT**
Telmatobius thompsoni **EN**
Telmatobius truebae **EN**
Telmatobius verrucosus **VU**
Telmatobius yuracare **VU**
Thoropa lutzi **EN**
Thoropa petropolitana **VU**
Pseudacris triseriata **LC**
Pseudophryne corroboree **CR**
Ptychohyla **CR**
Ptychohyla dendrophasma **CR**
Ptychohyla euthysanota **NT**
Ptychohyla leonhardschultzei **EN**
Ptychohyla panchoi **EN**
Ptychohyla sanctaecrucis **CR**
Ptychohyla zophodes **DD**
Rana sierrae **EN**
Rhacophorus margaritifer **LC**
Rheobatrachus vitellinus **EX**
Rhinella chrysophora **EN**
Rhinoderma rufum **CR**
Scinax heyeri **DD**
Smilisca cyanosticta **NT**
Strabomantis necerus **VU**
Taudactylus acutirostris **CR**
Taudactylus eungellensis **CR**
Taudactylus pleione **CR**
Telmatobius arequipensis **VU**
Telmatobius atahualpai **DD**
Telmatobius brevipes **EN**
Telmatobius carrillae **VU**
Telmatobius cirrhacelis **CR**
Telmatobius contrerasi **DD**
Telmatobius dankoi **DD**
Telmatobius edaphonastes **EN**
Telmatobius hauthali **VU**
Telmatobius hypselocephalus **EN**
Telmatobius intermedius **DD**
Telmatobius laticeps **EN**
Telmatobius marmoratus **VU**
Telmatobius necopinus **EN**
Telmatobius pefauri **CR**
Telmatobius philippii **DD**
Telmatobius pisanoi **EN**
Telmatobius schreiteri **EN**
Telmatobius sibiricus **EN**
Telmatobius stephani **EN**
Telmatobius timens **DD**
Telmatobius vellardi **CR**
Telmatobius vilamensis **DD**
Telmatobius zapahuirensis **CR**
Thoropa miliaris **LC**
Thoropa saxatilis **NT**

BIBLIOGRAPHY

55 references found for **Batrachochytrium dendrobatidis**

Managment information
Australian Department of the Environment and Heritage, 2004. Chytridiomycosis (Amphibian Chytrid Fungus Disease). Australia s Natural Heritage Trust.

Berger, Lee, Alex D. Hyatt, Veronica Olsen, Sandra G. Hengstberger Donna Boyle, Gerry Marantelli, Kaye Humphreys, Joyce E. Longcore., 2002. Production of polyclonal antibodies to *Batrachochytrium dendrobatidis* and their use in an immunoperoxidase test for chytridiomycosis in amphibians. Diseases of Aquatic Organisms Vol 48 213-220.
Summary: Available from: http://www.jcu.edu.au/school/phtm/PHTM/frogs/papers/berger-2002-polyab.pdf [Accessed 14 September 2005]
Berger, L., Hyatt, A.D., Olsen, V., Hengstberger, S.G., Boyle, D., Marantelli, G., Humphreys, K., Longcore, J.E. 2002. Production of Polyclonal Antibodies to *Batrachochytrium dendrobatidis* and Their use in an Immunoperoxidase Test for Chytridiomycosis in Amphibians, Dis Aquat Organ. 48 (3): 213 -220. (Abstract)
Berger, L., Speare, R., and Hyatt., A.D. 1999. Chytrid Fungi and Amphibian Declines: Overview, Implications and Future Directions. Declines and Disappearances of Australian Frogs. 23 - 34.
Berger, L., Speare, R., and Kent, A. 1999. Diagnosis of chytridiomycosis in amphibians by histologic examination.
Summary: This paper outlines techniques for identifying the chytrid fungus.
Available from: http://www.jcu.edu.au/school/phtm/PHTM/frogs/histo/chhisto.htm [Accessed 17 December 2004]
Daszak, P., A. Strieby, A. A. Cunningham, J. E. Longcore, C. C. Brown and D. Porter., 2004. Experimental Evidence That The Bullfrog (*Rana catesbeiana*) Is A Potential Carrier Of Chytridiomycosis, An Emerging Fungal Disease Of Amphibians. Herpetological Journal, Vol. 14, Pp. 201-207 (2004)
Summary: Available from: http://www.conservationmedicine.org/papers/Herp_J_Bullfrog_exptl_infection.pdf [Accessed 14 September 2005]
DeWeerdt , Sarah. 2001. Coordinating an International Monitoring Program The Declining Amphibian Populations Task Force. Conservation in Practice Winter 2001 Vol 2 no. 1
Garthwaite, R. Department of Conservation. *Batrachochytrium dendrobatidis*, Frog Chytrid Fungus. Department of Conservation: Waikato.
Gascon, C., Collins, J. P., Moore, R. D., Church, D. R., McKay, J. E. and Mendelson, J. R. III (eds). 2007. Amphibian Conservation Action Plan. IUCN/SSC Amphibian Specialist Group. Gland, Switzerland and Cambridge, UK. 64pp.
Summary: The Amphibian Conservation Action Plan (ACAP) is designed to provide guidance for implementing amphibian conservation and research initiatives at all scales from global down to local.
Available from: http://www.amphibians.org/newsletter/ACAP.pdf [Accessed 9 June 2008]
Johnson, Megan L & Richard Speare, 2005. Possible modes of dissemination of the amphibian chytrid *Batrachochytrium dendrobatidis* in the environment. Dis Aquat Org Vol. 65: 181-186, 2005
Summary: Available from: http://www.int-res.com/articles/dao2005/65/d065p181.pdf [Accessed 14 September 2005]
Johnson, M.L. and Speare, R. 2003. Survival of *Batrachochytrium dendrobatidis* in Water: Quarantine and Disease Control Implications, Emerging Infectious Diseases 9 (8): 922 - 925.
Summary: Available from: http://www.cdc.gov/ncidod/EID/vol9no8/03-0145.htm [Accessed 7 Dec 2004]
Johnson, M.L., Berger, L., Philips, L. and Speare, R. 2003. Fungicidal Effects of Chemical Disinfectants, UV Light, Desiccation and Heat on the Amphibian Chytrid *Batrachochytrium dendrobatidis*, Diseases of Aquatic Organisms 57 (3): 255 - 260.
Mazzoni, R., Cunningham, A.A., Daszak, P., Apolo, A. Perdomo, P. and Speranza., G. 2003. Emerging Pathogen of Wild Amphibians in Frogs (*Rana catesbeiana*) Farmed for International Trade, Emerging Infectious Diseases 9 (8): 995 - 998.
Michigan Frog Survey. 2003. Michigan Frog Survey Update. Michigan Department of Natural Resources Wildlife Division Natural Heritage Program.
Parker, J.M., Mikaelian, I., Hahn, N. and Diggs, H.E. 2002. Clinical Diagnosis and Treatment of Epidermal Chytridiomycosis in African Clawed Frogs (*Xenopus tropicalis*), Comp Med. 52 (3): 265 – 268. (Abstract)
Rollins-Smith, L., Reinert, L.K., Miera, V. and Conlon, J.M. 2002. Antimicrobial Peptide Defenses of the Tarahumara Frog, *Rana tarahumarae*, Biochemical and Biophysical Research Communications 297 (2): 361 - 367.
Ron, Santiago R., 2005. Predicting the Distribution of the Amphibian Pathogen *Batrachochytrium dendrobatidis* in the New World1. Biotropica 37 (2), 209-221.
Speare, R. and Core Working Group of Getting the Jump on Amphibian Disease. 2001. Nomination for Listing of Amphibian Chytridiomycosis as a key Threatening Process Under the Environment Protection and Biodiversity Conservation act 1999. In: Speare, R. and Steering Committee of Getting the Jump on Amphibian Disease. Developing Management Strategies to Control Amphibian Diseases: Decreasing the Risk due to Communicable Diseases. School of Public Health and Tropical Medicine, James Cook University: Townsville. 163 - 187.
Speare R, Berger L. Global distribution of chytridiomycosis in amphibians.
Summary: This document gives details on the global distribution of the chytrid fungus, and was last updated in April 2004.
Available from: http://www.jcu.edu.au/school/phtm/PHTM/frogs/chyglob.htm. [Accessed 11 November 2000].
USDI (United States Department of the Interior) U.S. Fish and Wildlife Service. 2003. RE: Buck Springs Range Management Allotment Plan
Van-Ells, T., Stanton, J., Strieby, A., Daszak, P., Hyatt, A.D. and Brown, C. 2003. Use of Immunohistochemistry to Diagnose Chytridiomycosis in Dyeing Poison Dart Frogs (*Dendrobates tinctorius*), Journal of Wildlife Diseases 39 (3): 742 - 745.

General information

Bell, Ben D., Scott Carver, Nicola J. Mitchell, Shirley Pledger., 2004. The recent decline of a New Zealand endemic: how and why did populations of Archey's frog *Leiopelma archeyi* crash over 1996-2001? Biological Conservation 120 (2004) 189-199
Summary: Available from: http://www.jcu.edu.au/school/phtm/PHTM/frogs/papers/bell-2004.pdf [Accessed 14 September 2005]
Berger, L, R Speare, HB Hines, G Marantelli, AD Hyatt, KR Mcdonald, LF Skerratt, V Olsen, JM Clarke, G Gillespie, M Mahony, N Sheppard, C Williams And MJ Tyler., 2004. Effect of season and temperature on mortality in amphibians due to chytridiomycosis. Australian Veterinary Journal Volume 82, No 7, July 2004
Summary: Available from: http://www.ava.com.au/avj/private/0407/04070434.pdf [Accessed 14 September 2005]
Berger, L., Speare, R., Daszak, P., Green, D.E., Cunningham, A.A., Goggin, C.L., Slocombe, R., Ragan, M.A., Hyatt, A.D., McDonald, K.R., Hines, H.B., Lips, K.R., Marantelli, G. and Parkes, H. 1998. Chytridiomycosis Causes Amphibian Mortality Associated With Population Declines in the Rain Forests of Australia and Central America, Population Biology (Proc Natl Acad Sci U S A.) 95 (15): 9031 – 9036.
Summary: Available from: http://www.pnas.org/cgi/content/full/95/15/9031 [Accessed 7 Dec 2004]

Berger, L., Speare, R., Hines, H.B., Marantelli, G., Hyatt, A.D., McDonald., K.R., Skerratt, L.F., Olsen, V., Clarke, J.M., Gillespie, G., Mahony, M., Sheppard, N. Williams, C. and Tyler. M.J. 2004. Effect of Season and Temperature on Mortality in Amphibians due to Chytridiomycosis, Australian Veterinary Journal 82 (7): 434 - 439.

Bosch, J., Martinez-Solano, I., and Garcia-Paris, M. 2000. Chytridiomycosis in Spain: First European Report of Declines of Wild Amphibians Associated with Chytridiomycosis.

Summary: This article gives details about the first caseof chytrid fungus in Spain.
Available from: http://www.jcu.edu.au/school/phtm/PHTM/frogs/spain/spainchy.htm [Accessed 17 December 2004]

Carey, C., Cohen, N. and Rollins-Smith, L. 1999. Amphibian declines: an immunological perspective. Developmental and Comparative Immunology. 23 (6): 459-472.

Summary: This paper discusses the role of disease in amphibian decline, and the immunological response.

Commonwealth Scientific and Industrial Research Organisation (CSIRO), 2003. Researching Frog Fungus.

Dasak, P., Andrew, A Cunningham and Hyatt, D Alex., 2003. Infectious disease and amphibian population declines. Diversity and Distributions 9, 141-150

Daszak, P., Berger, L., Cunningham, A.A., Hyatt, A.D., Green, D.E., Speare, R., 1999. Emerging Infectious Diseases and Amphibian Population Declines. Emerg Infect Dis [serial on the Internet]. November-December 1999

Summary: Available from: http://www.cdc.gov/ncidod/eid/vol5no6/daszak.htm [Accessed 7 Dec 2004]

Davidson, E.W., Parris, M., Collins, J.P., Longcore, J.E., Pessier, P.A. and Brunner, J. 2003. Pathogenicity and Transmission of Chytridiomycosis in Tiger Salamanders (*Ambystoma tigrinum*), Copeia 3: 601 - 607.

Fellers, G.M., Green, D.E. and Longcore, J.E. 2001. Oral Chytridiomycosis in the Mountain Yellow-legged Frog (*Rana muscosa*), Copeia 4: 945 - 953.

Hero, Jean-Marc & Clare Morrison., 2004. Frog declines in Australia Global implications. Herpetological Journal Vol. 14, pp. 175-186 (2004)

Summary: Available from: http://www.griffith.edu.au/school/asc/ppages/academic/jmhero/EndgFrogs/docs/Hero%20&%20Morrison%202004.pdf [Accessed 14 September 2005]

IUCN, Conservation International, and NatureServe. 2006. Global Amphibian Assessment. Downloaded on 4 May 2006.

Summary: The Global Amphibian Assessment (GAA) is the first-ever comprehensive assessment of the conservation status of the world s 5,918 known species of frogs, toads, salamanders, and caecilians. This website presents results of the assessments, including IUCN Red List threat category, range map, ecology information, and other data for every amphibian species.
Available from: http://www.globalamphibians.org/ [Accessed 5 November 2006].

Johnson, Pieter.T.J., 2006. Amphibian diversity: Decimation by dusease. Published online before print February 21, 2006, 10.1073/pnas.0600293103

Summary: Available from: http://www.pnas.org/cgi/content/full/103/9/3011 [Accessed 14 August 2006]

Kingsley D. Environment News, 23 April 2002.

Summary: This article gives details about the first reports of chytrid fungus in Archey s frog.
Available from: http://www.abc.net.au/science/news/enviro/EnviroRepublish_537533.htm [Accessed 17 December 2004]

Lips, Karen.R, Forrest Brem, Roberto Brenes, John D. Reeve, Ross A. Alford, Jamie Voyles, Cynthia Carey, Lauren Livo, Allan P. Pessier, and James P. Collins., 2006. Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. Published online before print February 15, 2006, 10.1073/pnas.0506889103

Summary: Available from: http://www.pnas.org/cgi/content/full/103/9/3165 [Accessed 14 August 2006] Available from: http://www.pnas.org/cgi/content/full/103/9/3165 [Accessed 14 August 2006]

Lips, K.R., Green, D.E. and Papendick, R. 2003. Chytridiomycosis in Wild Frogs from Southern Costa Rica, Journal of Herpetology 37 (1): 215 - 218.

Lips, K.R., Mendelson, J.R. Munoz-Alonso, A., Canseco-Marquez, L. and Mulcahy, D.G. 2004. Amphibian Population Declines in Montane Southern Mexico: Resurveys of Historical Localities, Biological-Conservation 119 (4): 555 - 564.

Morehouse, E.A., James, T.Y., Ganley, A.R.D., Vilgalys, R., Berger, L., Murphy , P.J. and Longcore, J.E. 2003. Multilocus Sequence Typing Suggests the Chytrid Pathogen of Amphibians is a Recently Emerged Clone, Molecular Ecology 12 (2): 395 - 403.

Muths, E., Corn, P.S., Pessier, A.P. and Green, D.E. 2003. Evidence for Disease-related Amphibian Decline in Colorado, Biological Conservation 110 (3): 357 - 365.

Mutschmann, F., Berger, L., Zwart, P. and Gaedicke, C. 2000. Chytridiomycosis in Amphibians: First Report in Europe, Berl Munch Tierarztl Wochenschr 113 (10): 380 – 383. (Abstract)

Norman, R. Undated. Chytrid fungus disease in New Zealand. Massey University Institute of Veterinary, Animal and Biomedical Sciences.

Summary: Article outlining the first case of chytrid fungus in New Zealand.
Available from: http://wildlife.massey.ac.nz/research/chytrid/chytrid1.asp [Accessed 17 December 2004]

Parris, M. J. 2004. Hybrid response to pathogen infection in interspecific crosses between two amphibian species (Anura: Ranidae). Evolutionary Ecology Research 6: 457-471.

Summary: *B. dendrobatidis* differentially affects genotypes between two species of hybridizing leopard frogs (*Rana*). Hybrid genotypes are more susceptible to infection, and suffer greater reductions in growth and development from the fungus.

Parris, M.J. and Beaudoin, J.G. 2004. Chytridiomycosis Impacts Predator-prey Interactions in Larval Amphibian Communities, Oceologia (Berlin) 140 (4): 626 - 632.

Summary: *B. dendrobatidis* alters the outcome of natural predator - prey dynamics in a larval amphibian - predator system.

Parris, M. J. and D. R. Baud. 2004. Interactive effects of a heavy metal and chytridiomycosis on gray treefrog larvae (*Hyla chrysoscelis*). Copeia 2004: 343-349.

Summary: *B. dendrobatidis* impacts on *Hyla* larvae may be somewhat ameliorated in a heavy metal (Cu) aquatic environment. Thus, pathogenic effects may be a result of interactions with other aquatic contaminants.

Parris, M. J. and T. O. Cornelius. 2004. Fungal pathogen causes competitive and developmental stress in larval amphibian communities. Ecology 85: 3385-3395.
Summary: This paper documents that *B. dendrobatidis* induces competitive effects in the larval environment between a toad (*Bufo*) and treefrog (*Hyla*) species.
Rollins-Smith, L.A., Carey, C., Longcore, J., Doersam, J.K., Boutte, A., Bruzgal, J.E., and Conlon, J.M. 2002. Activity of antimicrobial skin peptides from ranid frogs against *Batrachochytrium dendrobatidis*, the chytrid fungus associated with global amphibian declines. Developmental and Comparative Immunology. 26 (5): 471-479.
Summary: This paper outlines the role of antimicrobial peptides in deterring chytrid infection.
Speare R, Berger L. Chytridiomycosis in amphibians in Australia.
Summary: Available from: http://www.jcu.edu.au/school/phtm/PHTM/frogs/chyspec.htm. [Accessed 9 October 2000] .
USGS (U.S. Geological Survey). 2000. Research Project: Review and Classification of Visitor Impacts to Wildlife Research Methods. U.S. Department of the Interior: Patuxent Wildlife Research Center.
Summary: Available from: http://www.pwrc.usgs.gov/research/sis2000/longco07.htm [Accessed 7 Dec 2004]
Voyles, Jamie., Sam Young, Lee Berger, Craig Campbell, Wyatt F. Voyles, Anuwat Dinudom, David Cook, Rebecca Webb, Ross A. Alford, Lee F. Skerratt, Rick Speare. 2009. Pathogenesis of Chytridiomycosis, a Cause of Catastrophic Amphibian Declines. Science, Vol. 326 No. 5952, October 23, 2009.
Summary: The pathogen *Batrachochytrium dendrobatidis* (Bd), which causes the skin disease chytridiomycosis, is one of the few highly virulent fungi in vertebrates and has been implicated in worldwide amphibian declines. However, the mechanism by which Bd causes death has not been determined. We show that Bd infection is associated with pathophysiological changes that lead to mortality in green tree frogs (*Litoria caerulea*). In diseased individuals, electrolyte transport across the epidermis was inhibited by >50%, plasma sodium and potassium concentrations were respectively reduced by ~20% and ~50%, and asystolic cardiac arrest resulted in death. Because the skin is critical in maintaining amphibian homeostasis, disruption to cutaneous function may be the mechanism by which Bd produces morbidity and mortality across a wide range of phylogenetically distant amphibian taxa
Waldman, B., van de Wolfshaar, K.E., Klena, J.D., Andjic, V., Bishop, P., and Norman, R. J. de B. 2001. Chytridiomycosis in New Zealand frogs. Surveillance. 28 (3): 9-11.
Summary: This article gives details about the first case of chytrid fungus in New Zealand, including possible means of introduction and spread.
Available from: http://ivabs.massey.ac.nz/centres/wildlife/rschrepts/chytrid/chytrid_article.pdf [Accessed 17 December 2004]
Weldon C, du Preez LH, Hyatt AD, Muller R, Speare R., 2004. Origin of the amphibian chytrid fungus. Emerg Infect Dis [serial on the Internet]. 2004 Dec.
Summary: Available from http://www.cdc.gov/ncidod/EID/vol10no12/03-0804.htm [Accessed 14 December 2005]
Woodhams, D.C., Alford, R.A. and Marantelli, G. 2003. Emerging Disease of Amphibians Cured by Elevated Body Temperature, Diseases of Aquatic Organisms 55 (1): 65 - 67.
Young, B.E., Lips, K.R., Reaser, J.K., Ibanez, R., Salas, A.W., Rogelio Cedeno, J., Coloma, L.A., Ron, S., La Marca, E., Meyer, J.R., Munoz, A., Bolanos, F., Chaves, G. and Romo, D. 2001. Population declines and priorities for amphibian conservation in Latin America. Conservation Biology. 15 (5): 1213-1223.
Summary: A discussion of the factors involved in the population declines of amphibians in Latin America.