

Bidens pilosa [简体中文](#) [正體中文](#)

System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Plantae	Magnoliophyta	Magnoliopsida	Asterales	Asteraceae

Common name

tombo-maga (English, Sierra Leone), kiradale (English, Ivory Coast), eyinata (English, Nigeria), nana (English, Sierra Leone), zikilli wissi (English), akesan (English, Nigeria), dwirantwi (English, Ghana), pilipili (English), klakuo (English), anansee mpaane (English, Ghana), sanyina (English, Sierra Leone), dzani pipi (English, Ghana), gyinantwi (English, Ghana), beggar's tick (English), kurofidie (English, Ghana), kete kete (English, Nigeria), cobbler's peg (English), fisi'uli (Tongan), kichoma nguo (Swahili), kichoma mguu (Swahili), Spanish needle (English), tabason (English, Ivory Coast), asta de cabra (Spanish), kandane (English, Sierra Leone), acetillo (Spanish), passoklo (English, Ivory Coast), amonoablanfè (English, Ivory Coast), kukwe kwo (English, Ivory Coast), Zweizhan (German), sornet (French), lebason (English, Ivory Coast), broom stick (English), hairy beggar ticks (English), piquants noirs (French), black fellows (English), black jack (English), devil's needles (English), pisau-pisau (English), herbe villebague (French), matua kamate (Fijian), broom stuff (English), batimadramadramatakaro (Fijian), herbe d'aiguille (French), mbatikalawau (Fijian), mbatimandramandra (Fijian), bident poilu (French), dada (English, Sierra Leone), carrapicho-deagulha (Portuguese), dadayem (Ibatan), nehe (Hawaiian), ki (Hawaiian), ki pipili (Hawaiian), sosolé (English, Ivory Coast), ki nehe (Hawaiian), alonga (English, Ivory Coast), sirvulaca (Spanish), arponcito (Spanish), aseduro (English, Ghana), kofetonga (Niuean), amor seco (Spanish), pipiripi (Maori), ko-sendagusa (Japanese), niroa (Maori), kamik tuarongo (Maori), kofetoga (Niuean), tebasson (English, Ivory Coast), pipiripi niroa (Maori), pipiripi kerekere (Maori), sanyi (English, Sierra Leone), nidul-lif (English, Sierra Leone), masquia (Spanish), bidente pilosa (Spanish), agberi-oku (English, Sierra Leone), adzrskpi (English, Ivory Coast), nangua (English, Ivory Coast), cadillo (Spanish), cacha de cabra (Spanish), légué (English, Ivory Coast), zebeyuzébogue (English, Ivory Coast), hierba amarilla (Spanish), diandu (English, Ivory Coast), rosilla (Spanish), dinenkui (English, Ivory Coast), nanguadian (English, Ivory Coast), kokosa (English, Ivory Coast), mazote (Spanish), diaani (English, Ivory Coast), iuna (English, Ivory Coast), perca (Spanish), gonoretti (English, Ivory Coast), niani (English, Liberia), zagoi ini (English, Ivory Coast), tagiaani (English, Ivory Coast), manamendigo (English, Ivory Coast), abissawa (English, Ivory Coast), puriket (English), pega-prga (Spanish), nguad (English), papunga chipaca (Spanish), alongoï (English, Ivory Coast), anasipagné (English, Ivory Coast), bident hérissé (French), iréné (English, Ivory Coast), picão-preto (Portuguese), pétéoré (English, Ivory Coast), zagaï zagagbé (English, Ivory Coast), zegbei zegbagwè (English, Ivory Coast)

Synonym

Bidens leucantha , (L.) Willd.
Bidens leucantha , Willd. var. *sundaica* (Blume) Hassk.
Bidens sundaica , (Blume)
Coreopsis leucantha , L.
Bidens odorata

Similar species

Summary

Bidens pilosa is a cosmopolitan, annual herb which originates from tropical and Central America. Its hardiness, explosive reproductive potential, and ability to thrive in almost any environment have enabled it to establish throughout the world. Generally introduced unintentionally through agriculture or sometimes intentionally for ornamental purposes, *B. pilosa* is a major crop weed, threat to native fauna, and a physical nuisance.



[view this species on IUCN Red List](#)

Species Description

Bidens pilosa is an erect, annual herb which stands from 0.3-2 m high and bears opposite, pinnately compound, broadly ovate, (3-)5-9-lobed leaves 3-20 cm long and 2.5-12 cm wide. Leaf segments ovate to lanceolate lobed or bilobed at the base with margins crenate-serrate and apices acute. Stems are reddish tinged; 4-angled, simple, or branched. Heads solitary or in lax paniculate cymes at the ends of the main stem and lateral branches, usually radiate, 5 – 12 mm broad. Heads with 2 rows of involucre bracts, outer ones 7-10, spatulate, reflexed at anthesis, 3-4 mm long, inner ones ovate lanceolate; ray flowers absent or 4-8, sterile, corolla 7-15 mm long, white to yellow or pinkish, disk flowers with 3.5 – 5 mm long, yellow corolla. Achenes are black, 4-8 ribbed, linear, 6-16 mm long, with 2-3(-5) retrorsely barbed bristles of 2-4 mm long (Aluka, undated; PIER 2007).

Lifecycle Stages

Bidens pilosa grows quickly. Plants flower 4 months after germination and produce mature seeds 4 weeks after flowering. Plants typically bear 80 flower heads with seeds with potential production of 3000 plants in a generation and 4 generations per year (DPI, 2008; Mvere, 2004; PIER, 2007).

Uses

Bidens pilosa is used as a medicinal plant in areas of Africa, Asia, and tropical America. Its roots, leaves, and seeds are reported to have antibacterial, antidiarrheal, anti-inflammatory, antimicrobial, antimalarial, diuretic, hepatoprotective, and hypotensive properties. In Africa, *B. pilosa* is used to treat headaches, ear infections, hangovers, diarrhoea, kidney problems, malaria, jaundice, dysentery, burns, arthritis, ulcers, and abdominal problems. It is also used as an anaesthetic, coagulant, and treatment to ease child birth. In sub-Saharan Africa, its fresh or dried shoots and young leaves are eaten as a leaf vegetable, especially in times of food scarcity. *B. pilosa* is also an ingredient of sauces eaten with many staple foods there (Mvere, 2004).

Habitat Description

Bidens pilosa is a hardy weed capable of invading a vast range of habitats ranging from moist soil, sand, limerock, or dry, infertile soil and low to high altitudes of up to 3,600 m. It thrives in disturbed areas, high sunlight, and moderately dry soils, but is known to invade grassland, heathland, forest clearings, wetlands, plantations, streamlines, roadsides, pasture, coastal areas, and agriculture areas. *B. pilosa* is capable of surviving severe droughts with a required annual rainfall range is 500-3500 mm. It is tolerant to a pH range of 4-9 and high salinities of up to 100 mM NaCl. It prefers temperatures above 15°C and below 45°C but is tolerant to frosts with roots capable of withstanding and regenerating after temperatures as low as -15°C. *B. pilosa* is not fire tolerant but is known to quickly invade burnt areas (PIER, 2007; Aluka, undated; DPI, 2008).

Reproduction

Sexual by self or cross-pollination. A single plant may produce 3,000-6,000 seeds per year which are spread by attaching to animals, birds, and people or dispersal by wind and water. Its full reproductive cycle may be completed in 57-70 days and be completed 5-6 times a years in some areas. Seeds are reported to have no dormancy, remain viable for 5-6 years, and a 74% germination rate in the field (PIER, 2007; Zungsontiporn, undated; DPI, 2008)

General Impacts

Bidens pilosa is a problematic species for many reasons throughout its range. A troublesome weed to at least 30 crops in over 40 countries, *B. pilosa* is known to significantly reduce crop yields. One study found that dry bean, *Phaseolus vulgaris*, harvests were reduced by 48% in Uganda and 18-48% in Peru due to impacts by *B. pilosa*. It forms dense stands that can out compete, out grow, and eliminate crop and native vegetation, specifically the lower vegetative strata, over large areas. *B. pilosa* prevents the regeneration of these plants as well, given its allelopathic properties. Leaf and root extracts are known to significantly suppress germination and seedling growth of many plants and are believed to remain active throughout decomposition. Furthermore, *B. pilosa* grows three times faster than similar plant species. All of these properties render it a quite formidable competitor.

Its thick stands impede access to roads, trails, and recreational areas, are a nuisance to travellers and tourists, and inflict damage to pavements and walls. Its burrs are a nuisance to people, as well as, sheep and other fleece producing livestock. The burrs are also a troublesome seed contaminant as they are difficult to separate. *Bidens pilosa* is also a host and vector to harmful parasites such as Root knot nematodes (*Meloidogyne* sp.) and Tomato spotted wilt virus (*Sclerotinia sclerotiorum*) (DPI, 2008; Mvere, 2004).

Management Info

Physical: *Bidens pilosa* is susceptible to hand weeding. Germination may be prevented by mulches if they are thick enough (PIER, 2007).

Chemical: *B. pilosa* is susceptible to several types of herbicides. Residual herbicides: diuron, bromacil, atrazine, simazine, ropazine, hexazinone, oryzalin, and ametryn; translocated herbicides: 2,4-D, glyphosate, amitrole, metribuzin, and dicamba; and contact herbicides bentazone, diquat, and paraquat have all been evaluated as effective means of controlling *B. pilosa* when applied at standard rates. *B. pilosa* is thought susceptible to the majority of broad-leafed plant herbicides (PIER, 2007).

Pathway

Bidens pilosa has been introduced to many new locations by man for agricultural or ornamental purposes (Carlquist, 1966).

Principal source: [Pacific Island Ecosystems at Risk \(PIER\)](#), 2007. *Bidens pilosa* L., Asteraceae

Compiler: National Biological Information Infrastructure (NBII) & IUCN/SSC Invasive Species Specialist Group (ISSG)

Review: Johan van Valkenburg, Dutch Plant Protection Service.

Publication date: 2010-08-30

ALIEN RANGE

[3] AMERICAN SAMOA	[7] AUSTRALIA
[1] AUSTRIA	[1] BELGIUM
[1] BENIN	[1] BOTSWANA
[1] BRITISH INDIAN OCEAN TERRITORY	[1] CAMBODIA
[1] CAMEROON	[2] CANADA
[1] CHINA	[1] CHRISTMAS ISLAND
[1] CONGO, THE DEMOCRATIC REPUBLIC OF THE	[13] COOK ISLANDS
[1] COTE D'IVOIRE	[1] CYPRUS
[1] CZECH REPUBLIC	[1] ECUADOR
[1] ESTONIA	[5] FIJI
[1] FRANCE	[16] FRENCH POLYNESIA
[1] GREECE	[1] GUAM

[1] INDONESIA	[1] ISRAEL
[1] ITALY	[1] JAPAN
[1] KENYA	[5] KIRIBATI
[1] LIBERIA	[1] MALAWI
[1] MALAYSIA	[2] MARSHALL ISLANDS
[1] MAURITIUS	[5] MICRONESIA, FEDERATED STATES OF
[1] MOZAMBIQUE	[7] NEW CALEDONIA
[3] NEW ZEALAND	[1] NIGERIA
[1] NIUE	[1] NORFOLK ISLAND
[2] NORTHERN MARIANA ISLANDS	[1] PALAU
[1] PAPUA NEW GUINEA	[1] PHILIPPINES
[1] PITCAIRN	[3] PORTUGAL
[2] SAMOA	[1] SEYCHELLES
[1] SIERRA LEONE	[1] SLOVENIA
[1] SPAIN	[1] TAIWAN
[1] TANZANIA, UNITED REPUBLIC OF	[1] THAILAND
[5] TONGA	[1] UGANDA
[3] UNITED KINGDOM	[26] UNITED STATES
[2] UNITED STATES MINOR OUTLYING ISLANDS	[1] VANUATU
[1] VIET NAM	[1] VIRGIN ISLANDS, U.S.
[2] WALLIS AND FUTUNA	[1] ZAMBIA
[1] ZIMBABWE	

BIBLIOGRAPHY

40 references found for *Bidens pilosa*

Management information

[Brandes, Dietmar., 2001. *Bidens pilosa* L. und ihre Einbürgerungschancen in den Ländern der Europäischen Union](#)

Summary: Available from:

http://bib1p1.rz.tu-bs.de/docportal/servlets/MCRFileNodeServlet/DocPortal_derivate_00001309/Document.pdf;jsessionid=0000w-cml6ZCGCzBvSIdELgEY6?hosts=local [Accessed on 7 July 2008]

Carlquist, Sherwin ., 1966. The Biota of Long-Distance Dispersal. II. Loss of Dispersibility in Pacific Compositae. *Evolution*, Vol. 20, No. 1 (Mar., 1966), pp. 30-48

[Department Of Primary Industries \(DPI\),. 2008. Victorian Resources Online Statewide. Cobblers Pegs \(*Bidens pilosa* L.\): Impact Assessment - Cobblers Pegs \(*Bidens pilosa* L.\) in Victoria](#)

Summary: Available from: http://hear.org/starr/publications/2006_lanai_islets_botanical_survey.pdf [Accessed on 7 July 2008]

[Department Of Primary Industries \(DPI\),. 2008. Victorian Resources Online Statewide. Cobblers Pegs \(*Bidens pilosa* L.\): Invasiveness Assessment - Cobblers Pegs \(*Bidens pilosa* L.\) in Victoria](#)

Summary: Available from: http://www.dpi.vic.gov.au/DPI/Vro/vrosite.nsf/pages/invasive_cobblers_pegs [Accessed on 7 July 2008]

[Department Of Primary Industries \(DPI\),. 2008. Victorian Resources Online Statewide. Cobblers Pegs \(*Bidens pilosa* L.\): Present distribution-Potential distribution](#)

Summary: Available from: http://www.dpi.vic.gov.au/DPI/Vro/vrosite.nsf/pages/weeds_herbs_annual_cobblers_pegs [Accessed on 7 July 2008]

[Heap, I., 2008. The International Survey of Herbicide Resistant Weeds. Group B/2 Resistant Hairy Beggersticks \(*Bidens pilosa*\) Brazil. Online. Internet. July 06, 2008 . Available \[www.weedscience.com\]\(http://www.weedscience.com\)](#)

Summary: Available from: <http://www.weedscience.org/Case/Case.asp?ResistID=42> [Accessed on 7 July 2008]

[Medeiros, A.C., L.L. Loope, and C.G. Chimera., 1993. Kanaio Natural Area Reserve Biological Inventory and Management Recommendations, March 1993, for the Natural Area Reserves System, State of Hawaii.](#)

Summary: Available from: <http://www.dofaw.net/nars/files/kanaioplan.doc> [Accessed on 7 July 2008]

[Mvere, B., 2004. *Bidens pilosa* L. In: Grubben, G.J.H. & Denton, O.A. \(Editors\). PROTA 2: Vegetables/Legumes. \[CD-Rom\]. PROTA, Wageningen, Netherlands.](#)

Summary: Available from: http://database.prota.org/PROTAhtml/Bidens%20pilosa_En.htm [Accessed on 7 July 2008]

[Space, James C., Barbara Waterhouse, Julie S. Denslow and Duane Nelson., 2000. Invasive Plant Species on Rota, Commonwealth of the Northern Mariana Islands. U.S.D.A. Forest Service Pacific Southwest Research Station Institute of Pacific Islands Forestry Honolulu, Hawaii i, USA](#)

Summary: Available from: http://hear.org/AlienSpeciesInHawaii/articles/pier/pier_rota_report.pdf [Accessed on 7 July 2008]

[Starr, Forrest; Kim Starr and Ken Wood., 2006. Lanai Offshore Islets Botanical Survey. Prepared for: Department of Land and Natural Resources, Division of Forestry and Wildlife and Offshore Islet Restoration Committee](#)

Summary: Available from: http://hear.org/starr/publications/2006_lanai_islets_botanical_survey.pdf [Accessed on 7 July 2008]

[Waterhouse, D.F. 1997. The major invertebrate pests and weeds of agriculture and plantation forestry in the southern and western Pacific. ACIAR Monograph No. 44, 99p.](#)



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Bidens pilosa*

General information

[Aluka, undated. *Bidens pilosa* L. \[family Compositae\]](#)

Summary: Available from:

<http://www.aluka.org/action/showCompilationPage?doi=10.5555/AL.AP.COMPILATION.PLANT-NAME-SPECIES.BIDENS.PILOSA&cookieSet=1> [Accessed on 7 July 2008]

Auld, Bruce., Hirohiko Morita, Tomoko Nishida, Misako Ito and Peter Michael., 2003. Shared exotica: Plant invasions of Japan and south eastern Australia. *Cunninghamia* (2003) 8(1): 147-152

[California: Information on California plants for education, research and conservation. \[web application\]. 2008. Taxon Report 1088 *Bidens pilosa* L. var. *pilosa*. Berkeley, California: The California Database \[a non-profit organization\].](#)

Summary: Available from: http://www.calflora.org/cgi-bin/species_query.cgi?where-calrecnum=1088 [Accessed on 7 July 2008]

Corlett, Richard T., 1988. The Naturalized Flora of Singapore *Journal of Biogeography*, Vol. 15, No. 4, Festschrift in Honour of Professor Donald Walker, (Jul., 1988), pp. 657-663

Corlett, Richard T., 1992. The Naturalized Flora of Hong Kong: A Comparison with Singapore. *Journal of Biogeography*, Vol. 19, No. 4 (Jul., 1992), pp. 421-430

Dafni, Amots and David Heller., 1982. Adventive flora of Israel ♦ Phytogeographical, ecological and agricultural aspects. *Plant Systematics and Evolution*. Volume 140, Number 1 / April, 1982

[DAISIE \(Delivering Alien Invasive Species Inventories for Europe\) 2008. *Bidens pilosa* L.](#)

Summary: Available from: <http://www.europe-aliens.org/speciesFactsheet.do?speciesId=122433> [Accessed on 7 July 2008]

[DAISIE \(Delivering Alien Invasive Species Inventories for Europe\) 2008. *Bidens pilosa* L. Distribution Map](#)

Summary: Available from: <http://www.europe-aliens.org/speciesFactsheet.do?speciesId=122433#> [Accessed on 7 July 2008]

Della, Athena & G. Iatrou., 1995. New Plant Records from Cyprus. *Kew Bulletin*, Vol. 50, No. 2 (1995), pp. 387-396

[Global Biodiversity Information Facility \(GBIF\), 2008. Species: *Bidens pilosa* L. Kofe Tonga](#)

Summary: Available from: <http://data.gbif.org/species/13746874> [Accessed 15 June 2010]

Groves, R.H. (Convenor), Hosking, J.R., Batianoff, G.N., Cooke, D.A., Cowie, I.D., Johnson, R.W., Keighery, G.J., Lepschi, B.J., Mitchell, A.A., Moerkerk, M., Randall, R.P., Rozefelds, A.C., Walsh N.G., and Waterhouse, B.M. 2003. Weed categories for natural and agricultural ecosystem management. Bureau of Rural Sciences, Canberra.

Summary: This document gives the status of all weed species found in Australia.

Hadac, Emil Z and Vera Hadacová., 1969. Notes on the ecology and distribution of *Bidens pilosa* L. in Cuba. *Folia Geobotanica* Volume 4, Number 2 / June, 1969

[Hyde, M.A. & Wursten, B. 2008. Flora of Zimbabwe: Species information: *Bidens pilosa* L.](#)

Summary: Available from: http://www.zimbabweflora.co.zw/speciesdata/species.php?species_id=160650 [Accessed on 7 July 2008]

[ITIS \(Integrated Taxonomic Information System\), 2008. Online Database. *Bidens pilosa* L.](#)

Summary: An online database that provides taxonomic information, common names, synonyms and geographical jurisdiction of a species. In addition links are provided to retrieve biological records and collection information from the Global Biodiversity Information Facility (GBIF) Data Portal and bioscience articles from BioOne journals.

Available from: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=35731 [Accessed on 7 July 2008]

[Leach, Helen., 2005. Gardens without weeds? Pre-European Maori gardens and inadvertent introductions. *New Zealand Journal of Botany*, 2005, Vol. 43: 271-284](#)

Summary: Available from: <http://www.rsnz.org/publish/nzjb/2005/012.pdf> [Accessed on 7 July 2008]

[Mattiske Consulting Pty Ltd., 2005. Flora and Vegetation on the Cloud Break and White Knight Leases. Prepared for: Fortescue Metals Group Limited \(FMG0501/29/05\)](#)

Summary: Available from: http://www.epa.wa.gov.au/docs/cloudbreak/PER_CloudBreak_AppD.pdf [Accessed on 7 July 2008]

McMullen, Conley K., 1987. Breeding Systems of Selected Galapagos Islands Angiosperms. *American Journal of Botany*, Vol. 74, No. 11, (Nov., 1987), pp. 1694-1705

Muller, Norbert and Shigetoshi Okuda., 1998. Invasion of Alien Plants in Floodplains-A comparison of Europe and Japan., *Plant Invasions: Ecological Mechanisms and Human Responses*, pp. 32 1-332 edited by U. Starfinger, K. Edwards, I. Kowarik and M. Williamson

Summary: Available from: http://hear.org/starr/publications/2006_lanai_islets_botanical_survey.pdf [Accessed on 7 July 2008]

New Zealand Plant Conservation Network., 2005. *Bidens pilosa* L. Weed Status

Summary: Available from: http://www.nzpcn.org.nz/exotic_plant_life_and_weeds/detail.asp?WeedID=1129 [Accessed on 7 July 2008]

[Pacific Island Ecosystems at Risk \(PIER\), 2007. *Bidens pilosa* L., Asteraceae](#)

Summary: Available from: http://www.hear.org/pier/species/bidens_pilosa.htm [Accessed on 7 July 2008]

Pattison, R.R; G. Goldstein & A. Ares., 1998. Growth, biomass allocation and photosynthesis of invasive and native Hawaiian rainforest species

[Plants of Hawaii, 2008. Images by Forest & Kim Starr *Bidens pilosa* Spanish needle, ki, ki nehe, ki pipili, nehe \(Asteraceae\)](#)

Summary: Available from: http://www.hear.org/starr/hiplants/images/thumbnails/html/bidens_pilosa.htm [Accessed on 7 July 2008]

Sherff, Earl E., 1916. Studies in the Genus *Bidens*. III. *Botanical Gazette*, Vol. 61, No. 6 (Jun., 1916), pp. 495-506

[Space, James C. and Marjorie Falanruw., 1999. Observations on invasive plant species in Micronesia. Prepared for the meeting of the Pacific Islands Committee, Council of Western State Foresters, Majuro, Republic of the Marshall Islands, February 22-26, 1999.](#)

Summary: Available from: <http://202.4.49.29/att/IRC/eCOPIES/INVASIVE%20SPECIES/micronesia.pdf> [Accessed on 7 July 2008]

[University of Hawaii Botany Department, 1998. Hawaiian Alien Plant Studies *Bidens pilosa* L. beggar s tick, Spanish needle Asteraceae](#)

Summary: Available from: http://www.botany.hawaii.edu/faculty/cw_smith/bid_pil.htm [Accessed on 7 July 2008]

[USDA, ARS, 1994. *Bidens pilosa* National Genetic Resources Program. Germplasm Resources Information Network - \(GRIN\) \[Online Database\]. National Germplasm Resources Laboratory, Beltsville, Maryland.](#)

Summary: Available from: <http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?7169> [Accessed on 7 July 2008]

[USDA, NRCS, 2008. PLANTS Database, Plants Profile for *Bidens pilosa* L. hairy beggarticks](#)

Summary: Available from: <http://plants.usda.gov/java/profile?symbol=BIPI> [Accessed on 7 July 2008]



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Bidens pilosa*

[Wu, Shan-Huah and Hsiang-Hua Wang., 2005. Potential Asteraceae Invaders in Taiwan: Insights from the Flora and Herbarium Records of Casual and Naturalized Alien Species. *Taiwania*, 50\(1\): 62-70, 2005](#)

Summary: Available from: <http://www.press.ntu.edu.tw/ejournal/Files/taiwan/200503/7.pdf> [Accessed on 7 July 2008]

[Zungsontiporn, Siriporn., undated. Invasive Alien Weeds in Thailand and Case Study on *Bidens pilosa* L. var *radiata* Shultz-Bip. Plant Protection Research and Development Office, Department of Agriculture, Jatujak, Bangkok, Thailand.](#)

Summary: Available from: <http://www.baphiq.gov.tw/public/Attachment/691512344971.pdf> [Accessed on 7 July 2008]