



Melaleuca quinquenervia  [简体中文](#)

System: Terrestrial

[正體中文](#)

Kingdom	Phylum	Class	Order	Family
Plantae	Magnoliophyta	Magnoliopsida	Myrtales	Myrtaceae

Common name melaleuca (English, Puerto Rico), cajeput (English), Mao-Holzrose (German), paper bark tree (English), five-veined paperbark tree (English), punk tree (English), itahou (English), niaouli (English, New Caledonia), capeputi (English), kinindrano (English, Madagascar), balsamo de cayeput (English), oli (English, Madagascar), aceite de cayeput (English), ahambo (English, Madagascar), white bottlebrush tree (English), paperbark teatree (English), belbowrie (English), niaouli (French), Japanese paper wasp (English), broad-leaved paperbark tree (English), corcho (English), numbah (English), kayu putih (English), bottle brush tree (English), broadleaf teatree (English), broadleaf paperbark tree (English)

Synonym *Metrosideros quinquenervia* , Cav.
Metrosideros coriacea , Poir.
Melaleuca leucadendron , var. *coriacea* Poir.
Melaleuca leucadendron ,  *angustifolia* L.f.
Melaleuca viridiflora , var. *angustifolia* L.f.
Melaleuca viridiflora , var.  *rubriflora* Brongn. & Gris
Melaleuca rubriflora , Vieillard ex Brongn. & Gris
Melaleuca leucadendron , var. *rubriflora* Brongn. & Gris
Melaleuca maideni , R.T. Baker
Melaleuca smithii , R.T. Baker
Melaleuca leucadendron , var. *albida*

Similar species

Summary The broad-leaved paperbark tree or melaleuca (*Melaleuca quinquenervia*) can reach heights of 25 meters and hold up to 9 million viable seeds in a massive canopy-held seed bank. This fire-resistant wetland-invader aggressively displaces native sawgrass and pine communities in south Florida, alters soil chemistry and modifies Everglades ecosystem processes. Melaleuca is notoriously difficult to control, however, bio-control (integrated with herbicidal and other methods) holds a promising alternative to traditional control methods.



[view this species on IUCN Red List](#)

Species Description

The Australian broad-leaved paperbark tree (herein referred to as "melaleuca") is a member of the Myrtaceae family, which comprises about 4000 to 5000 tropical and temperate species (Watson & Dallwitz 1992, in Dray Bennett & Center 2006). Recent research incorporating DNA studies has resulted in an improved classification of the family (Wilson *et al.* 2005). Earlier classifications in which a primary division was based on the possession of dry or fleshy fruit are not congruent with the evolutionary relationships shown by analysis of DNA sequence data. *Melaleuca*, which now includes *Callistemon*, belongs to the tribe Melaleuceae (Craven 2006, Wilson *et al.* 2005).

Melaleuca can reach 25 meters in height and grow to 90 centimeters in diameter. It is easily recognised by its spongy flaking bark (white, cream, orange-cream, fawn-grey or dark grey in colour (Craven, in press), lanceolate five-veined leaves, and clusters of woody seed capsules (Laroche 1999). Its white papery bark resembles birch and its white flower clusters resemble bottlebrush (Gioeli & Neal 2004). Its white tufted inflorescences are indeterminate, two to five centimeters long and arranged in bottlebrush-like spikes (Holliday 1989, in Center *et al.* 2006). Flowers of *M. quinquenervia*, like most Myrtaceae, have numerous stamens on a cup-shaped hypanthium surrounding the ovary. Myrtaceae leaves are simple and entire and the plants are usually aromatic (Laroche 1999); some have an intense citrus-like or eucalyptus oil odour when crushed (FLEPPC Undated). Within the spike-like inflorescence, flowers are clustered in threes; and secrete nectar, which collects within the base of the hypanthium; the stamens are arranged in five bundles each bundle consisting of five to ten fused stamens; the petals and filaments are usually white or creamish (Laroche 1999, L. Craven, pers. comm.). The capsular fruits, 2.7-4mm long, may persist for several years (Meskimen 1962, in Center *et al.* 2006). Please see the *Melaleuca Management Plan 1999* for botanical illustrations of *M. quinquenervia* (page 10).

Notes

M. quinquenervia is a member of the Myrtaceae (myrtle family) which also includes the *Eucalyptus* (gum) genus (Laroche 1999). *Melaleuca* (Myrtaceae) is the second largest genus in the Myrtaceae family and is represented by up to 250 species (Barlow 1986, in Turner *et al.* 1998), including a number of undescribed species.

M. quinquenervia is part of the broad-leaved *Melaleuca leucadendra*-complex, which contains 15 species that are endemic to the Australian-Tasmanian region (Craven 1999, in Wineriter *et al.* 2003). The name *Melaleuca* comes from the Greek, meaning black and white, presumably referring to the white bark, often charred black by fire (Debenham, 1962, in Turner *et al.* 1998).

Lifecycle Stages

Melaleuca trees may reach 90 years and still remain fertile (Serbesoff-King 2003).

Seedling and Sapling Stages: Seedlings appear to be less tolerant of harsh environmental conditions than are the seeds (Woodall 1983, in Turner *et al.* 1998). *Melaleuca* seeds germinate upon moist soils, usually within a few days of wetting, and may remain viable up to six months under water or in wet soils (Meskimen 1962, Myers 1975, in Laroche 1999). Seeds may germinate while completely inundated (Lockhart 1995, in Laroche 1999). Meskimen (1962, in Laroche 1999) found a trend for germination to occur more in sun than in shade. Rarity of seedlings within dense stands of *melaleuca* may be from either shading or allelopathic effects of *melaleuca* litter (DiStefano & Fisher 1983, in Laroche 1999). Seedlings less than several weeks or months old may die from fire or if soils are dry (Myers 1975, 1983) Droughts severe enough to lower the water table by one meter will also kill the seedlings (Woodall 1981a, in Turner *et al.* 1998). Seedlings are also less tolerant of fires, as they have a thinner, insulating bark layer (Woodall 1981a, in Turner *et al.* 1998). Soon after seedlings are able to withstand extreme conditions ranging from fire to total immersion for months (Meskimen 1962, in Laroche 1999). Young saplings and seedlings respond to inundation by changing leaf shapes. Leaves become more linear when meristems are deeply flooded, and more rounded when the meristem is nearer to the water surface (Laroche 1999). This adaptation may enable better light or nutrient utilisation, or help the saplings to survive flooding (Lockhart, 1996).



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Melaleuca quinquenervia*

Uses

Worldwide, many of the 4000-5000 Myrtaceae species are cultivated as ornamentals or as sources of fruits, spices, aromatic oils or timber (Laroche 1999). The thick, spongy bark has historically been used as fruit-packing, bedding material and insulation (von Mueller 1888, Morton 1966, in Dray Bennett & Center 2006). Ornamental/landscaping: *Melaleuca* spp. are often planted as ornamentals, for screening, for their interesting bark and for their showy flowers (Turner *et al.* 1998). The small crown and distinctive bark have made it a popular ornamental tree (Greary Undated). It is widely cultivated for erosion control, windbreaks and watershed cover (Little & Skomen 1989, in Munger 2005).

Wood products: The medium-density wood is difficult to season and tends to warp, but it finishes well as a cabinet wood (Greary Undated). Without preservative treatment it makes a poor fence post and a major deterrent to use is the high bark-to-wood ratio (Greary Undated). *Melaleuca* has been used extensively for carpentry and joinery work and is used for structural timber, fuel, pulpwood and insulation/stuffing and for traditional dwellings in its native New Caledonia. The bark is useful for its insulating properties and as a mulch and potting medium (Greary Undated; Brown & Duke 2000, in Munger 2005). Cutting and chipping operations are currently utilising *melaleuca* wood for landscape mulching and boiler fuel in Florida (Stocker 1999).

Honey-making: In Florida, the abundant flowering crop has been important to the apiary industry to sustain bee colonies and as a source of honey (Greary Undated). While *melaleuca* is believed to be an important component of Florida's beekeeping industry (a source of nectar for honey, package bees, and wax) there are no indications that flowers are a limiting factor for bees (Diamond *et al.* 1991, in Laroche 1999).

Essential oils: Essential oils are extracted from its leaves, twigs and seeds by hydrodistillation from plantations in New Caledonia (Doran & Turnball 1997, Doran 1999 in Ireland 2002) and Madagascar (Ramanoelina *et al.* 2008). Essential oils constitute a principal antiseptic component in some commercial disinfectants (Dray *et al.* 2006).

Habitat Description

In its native range melaleuca occurs in seasonally and permanently inundated wetlands along the eastern coast of Queensland and New South Wales, Australia (11°S to 34°S) (Holliday 1989, in Burrows & Balcunas 1997; Boland *et al.* 1987, in Center *et al.* 2006). Australian habitats that support melaleuca populations include low-lying coastal wetlands behind heath-dominated headlands, riparian zones, brackish estuaries, mangrove swamps (Rayamajhi *et al.* 2002), *Melaleuca* swamp forest, monsoon scrub, littoral rainforest, grassland, open forest, low shrubland on coastal dunes and lagoon margins (Craven, In press). In its invaded territories, melaleuca can infest relatively drier areas (Buckingham 2000) and invades a variety of forested and non-forested natural communities, including: freshwater marshes, wet grasslands, sawgrass prairies, disturbed cypress forests, wet pine flatwoods, Miami rock ridge pinelands, longleaf-slash pine, hardwood hammocks, salt marshes and mangroves.

In general, xeric communities such as scrub tend to be resistant, but not immune, to melaleuca invasion (Laroche 1999). Favourable moisture conditions are found in pine flatwood depressions and the broad ecotones where pine and dwarf pond cypress mix (Duever *et al.* 1986, in Munger 2005). Melaleuca is tolerant of fire, seasonal drought and seasonal flooding (see Gomes & Kozlowski 1980; Geary & Woodall 1990). Melaleuca can grow in sites that are nutrient-poor such as pine savannas or wet prairies (Woodall 1981) due to its ability to send vertical roots straight down to the water table (Munger 2005).

As observed in Florida, Pratt (2005b) suggests wetlands that experience moderate to short hydroperiods are the most vulnerable to invasion by melaleuca. Melaleuca invades disturbed land such as abandoned farmlands, depressions in stump-harvested pinelands, road/canal wetland construction sites, improved pasture, natural rangeland and urban areas (Duever *et al.* 1986, Myers 1983 1984, in Munger 2005). Undisturbed ecosystems can be resistant, but not immune to, melaleuca invasion (Ewel *et al.* 1976, in Laroche 1999); however, in south Florida melaleuca has invaded essentially every existing community (Laroche 1999).

In Australia melaleuca occurs on sand, sandy loam, sandstone, laterite over sand, silty soil and serpentine substrates (Craven, In press), in New Guinea on highly organic, alluvial clays and in New Caledonia on well-drained slopes, ridges in the uplands (Geary Undated) and on flat, poorly drained soils (L. Craven, pers. comm.). Melaleuca establishes best on sandy soils but it can survive on nearly any soil type in south Florida (Ewel 1986, Hofstetter 1991, in Munger 2005). It is commonly found in Everglades ecosystems characterised by high organic soils (Pratt *et al.* 2004) or limestone-derived soils (Geary & Woodall 1990). Although melaleuca is found in soils of high pH plants may perform better in slightly acidic soils (Kaufman 1999, in Munger 2005). Melaleuca in Hawaii grow well on calcareous beach sand and on soils derived from basalt ash and lava rock of pH 4.5-5.5 (Geary 1998, in Geary Undated). According to Woodall (1981) a map of soil pH cannot be used to predict melaleuca invasion.

In its native habitat melaleuca is found mainly from sea level to 100m, but occasionally at elevations of 1000m (Geary Undated). Most of southern Florida, where melaleuca readily invades, is less than 8m above sea level (Geary & Woodall 1990). In its native habitat mean annual rainfall ranges from 900-1250 mm; mean monthly temperatures range from 5°C -32°C and in the southernmost part of its range, a few light frosts occur per year (Geary Undated). Where frequent freezing temperatures become common, melaleuca becomes less invasive (Munger 2005). The tree grows successfully in its introduced range where rainfall is 5000mm and a winter maximum occurs (Geary 1998, in Geary Undated).

Reproduction

Melaleuca trees may flower within three years of germination, sometimes in the first year, and produce seed as many as five times per year (Meskimen 1962, in Laroche 1999; Meskimen 1962, in Turner *et al.* 1998). In Florida, synchronised flowering events occur during winter (from November to January) and to a lesser degree in the summer (although a small proportion may reproduce at non-synchronised intervals) (Meskimen 1962). Bursts of vegetative growth generally occurring after flowering (between January and February) (Laroche 1999). In wet years, flowering and growth can be extended from July to April to with several flowering cycles (Laroche 1999). The number of capsular fruit produced per centimeter of infructescence is greater among populations in it's introduced range (eg: Puerto Rico and Florida) than in it's native range (Pratt *et al.* 2005b; Pratt *et al.* 2007). Reproduction occurs along flower-bearing branch segments; persistent capsular fruits arise from flowers and are arranged in clusters, which may remain attached to the trunks, branches or twigs for up to ten years (Laroche 1999; Meskimen 1962, in Center *et al.* 2006). Melaleuca has two reproduction possibilities due to the fact that seed retention extends beyond seed ripening; first, a low-level, virtually continuous seed release ensures that at least some of the seeds on the ground near the tree will be fresh, thus allowing the species to exploit all reproduction opportunities no matter how short they are in duration; second, retention of seeds allows for a potential mass seed release if some natural catastrophe kills the tree (Woodall 1983, in Laroche 1999). Melaleuca is an extremely prolific seed producer. Capsules each contain 200 to 350 minute seeds (Meskimen 1962), and the canopy of a 21 meter high tree may produce 34 kg of mature capsules that contain up to 100 million seeds (Rayamajhi *et al.* 2002b, in Serbesoff-King 2003; Van Rayamajhi & Center 2005). Studies have shown that of these 10% to 15% contain embryos and of these embryonic seeds 62% are viable (Rayachhetry *et al.* 1998, in Serbesoff-King 2003), giving a total potential 9 million viable seeds per mature large tree; and one hectare may store as many as 25 billion seeds; this represents a massive canopy-held seed bank (Rayamajhi Unpub. Data, in Center *et al.* 2006). Results from seed burial tests indicate that seed viability is decreased by about 50% after eight months in the soil (Laroche 1999). Most buried seeds lose viability after about 1.5 years at seasonally flooded and permanently flooded sites, whereas seeds buried at non-flooded sites survived over a period up to 2 to 2.3 years (Van, Rayamajhi and Center 2005). The seed capsule must be dry before the seed will be released; anything that disrupts vascular connections thereby causing the capsules to desiccate and open will stimulate melaleuca seed release (Center *et al.* 2006). Desiccation can be caused by stem growth, cutting or breaking of the stem, fire damage, frost damage, self pruning due to shade, natural death of the tree or herbicide application (Laroche 1999; Woodall 1982, in Munger 2005). Melaleuca trees resprout and coppice readily (Conde *et al.* 1981, Hofstetter 1991, in Serbesoff-King 2003) following man-made cutting or natural damage. Extensive rooting and sprouting can occur from a fallen or cut tree (Serbesoff-King 2003). Melaleuca logs used as fence-posts frequently sprout new growth (Laroche 1999; Munger 2005). Melaleuca is pollinated by a variety of insects, most commonly honeybees, however, seed fertility is low. Pollination within the same flower results in reduced fruit set compared with pollination between flowers, promoting out-crossing. Flowering and seed production are less on shaded branches than on emergent canopy branches (Meskimen 1962, in Munger 2005).



General Impacts

For a detailed account of the impacts of *M. quinquenervia* please read: [Melaleuca quinquenervia \(Broad-Leaved Paperbark\) Impacts Information](#). The information in this document is summarised below.

Melaleuca is the most problematic invasive plant species in Florida because of its wide distribution range, prolific seed production and potential impact on human health (Fuller 2005). Melaleuca threatens the preservation of critical wildlife habitat in southern Florida including in the Florida Everglades National Park. Despite control efforts melaleuca still occurred in around 170 000 hectares of southern Florida in 1997, representing 6% of the total region (Bodley & Van 1999, in Rayamajhi *et al.* 2007; Laroche 1999).

Ecosystem Change: Melaleuca threatens the integrity of subtropical freshwater ecosystem processes in Florida (Dray & Center 1994, in Lopez-Zamora Comerford & Muchovej 2004) by altering soil chemistry, reducing decomposition rates and modifying hydrology and fire regime. Melaleuca also reduces species biodiversity and alters species composition.

Reduction in Native Biodiversity: *Melaleuca* forests provide limited food and habitat value for native wildlife and can reduce indices of native species in Florida wetlands by as much as 80% (Dray *et al.* 2006; Bodley *et al.*, 1994, O'Hare & Dalrymple, 1997, in Dray *et al.* 2009; Porazinska Pratt & Giblin-Davis 2007). Decreases in diversity of native plant biodiversity have also been linked with melaleuca in the Bahamas.

Habitat Alteration: Melaleuca is contributing to significant habitat loss in the Everglades National Park by converting fire-maintained sawgrass communities into *Melaleuca* forest (Turner *et al.* 1998, in Munger 2005).

Displacement: Melaleuca displaces pond cypress (*Taxodium ascendens*) (Myers 1975 1983, Ewel 1986, in Rayamajhi *et al.* 2008b), slash pine (*Pinus elliotii*) and sawgrass (*Cladium jamaicensis*) (Bodley *et al.*, 1994, in Tipping *et al.* 2008).

Competition: Melaleuca is competitively superior to most native vegetation occurring in the Florida Everglades (Turner *et al.* 1998, in Pratt *et al.* 2005b). It is fire-adapted, herbivore-adapted and produces seeds and roots prolifically.

Inhibits the Growth of Other Species: Allelochemicals present in roots can have a detrimental effect on the soil biota (Porazinska Pratt & Giblin-Davis 2007).

Economic: Balciunas and Center (1991, in Serbesoff-King 2003) reported that by the year 2010, close to \$2 billion would be lost due to the melaleuca invasion in southern Florida. Financial losses included \$1 billion in tourism to the Everglades NP, \$250 million in tourism to the rest of south Florida, \$250 million in recreation, \$250 million due to fires, \$1 million in control efforts, \$10 million due to loss of endangered species and \$1 million to nursery growers.

Agricultural: In one study 18 economic arthropod pests were collected from *M. quinquenervia* (Costello *et al.* 2008).

Human Health: As melaleuca populations expand in southern Florida and the human population increases the risk of fire and loss of human life and property increases (Laroche 1999).

Modification of Hydrology: A stand of melaleuca may transpire more water than the sawgrass communities it replaces (Hofstetter 1991a, in Laroche 1999).

Modification of Fire Regime: Ground fires, high temperatures, rapid spread rates and abundant smoke, all present in burning melaleuca stands, present new risks for wildlife in the Everglades wetlands (Flowers 1991, in Laroche 1999).

Modification of Nutrient Regime: The rate of decomposition of melaleuca litter is slower than that of native plants (Van & Rayamajhi, Unpub. Data, in Rayamajhi *et al.* 2006b).



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Melaleuca quinquenervia*

Management Info

For a detailed account of management of *M. quinquenervia* please read: [Melaleuca quinquenervia \(Broad-leaved Paperbark\) Management Information](#). The information in this document is summarised below.

Current management methods for melaleuca include herbicides, manual removal of plants, prescribed fires and bio-control.

Preventative Measures: Preventative measures are the best form of weed control. Education on the potential threats posed by melaleuca on invaded ecosystems should be targeted at the nursery industry and the general public.

Monitoring and Mapping: Model projections suggest there is considerable scope for further invasion of melaleuca under current climate conditions, with the highest risk areas occurring in Southeast Asia, the Caribbean, South and Central America and the Gulf coast in southern USA.

Physical: Mechanical removal using heavy equipment is not appropriate in most natural areas because of disturbances to soils and non-target native vegetation; however, this method of control can be applied along canal and utility rights-of-way (Laroche 1999).

Physical: Physical methods also include the use of prescribed fire and of flooding. More information is needed on the timing of prescribed burning, and constraints to this method include impacts on non-target species, the triggering of mass seed release by trees and liability concerns (Turner et al. 1998).

Chemical/Herbicidal Control: Exotic woody vegetation is most frequently managed by herbicides (Laroche 1999). Hexazinone and tebuthiuron are most effective in the control of melaleuca (Laroche 1999), however, they are no longer allowed to be applied directly to water in Florida (Laroche 1998a, in Serbesoff-King 2003). Current chemical control recommendations for melaleuca include low volume applications of glyphosate for control of saplings, and aerial or individual stem (girdle) applications of imazapyr alone, or in combination with glyphosate for mature trees (Langeland and Stocker 1997, in Stocker 1999).

Biological control: The lack of a long-lived soil seed bank (Van et al. 2005, in Center et al. 2007) makes *M. quinquenervia* vulnerable to herbivore-mediated reductions in fitness and delays in reproductive maturation. As canopy-held seed banks continue to diminish over time (Pratt et al. 2005), seedling suppression is predicted to have long-term effects on plant density. Two bio-control agents, the Australian melaleuca snout weevil (*Oxyops vitiosa*) and the Australian melaleuca psyllid (*Boreioglycaspis melaleucae*), have been approved by the USDA for use against melaleuca (Cuba et al. 2003, Wineriter et al. 2003, in Gioeli & Neal 2004) and have been released in the field. Research is being conducted on at least six other potential bio-control agents, including leaf, stem tip, and flower bud feeders (Burrows & Balciunas 1997 1998, Turner et al. 1998, in Stocker 1999).

Legislative: Melaleuca is on both the United States' Federal Noxious Weed List and the Florida Prohibited Aquatic Plant List (Class I Prohibited aquatic plant) (Florida Department of Environmental Quality).

Integrated management: As a result of the implementation of the integrated *Melaleuca Management Plan 1999* almost 100 000 acres of natural area have been cleared of melaleuca (Laroche 1994).

[The Areawide Management Evaluation of Melaleuca quinquenervia \(TAME\)](#) aims to demonstrate the effectiveness of integrated control of melaleuca in invaded habitats in the United States and elsewhere.

Pathway

Melaleuca quinquenervia has been internationally disseminated over the course of the last century for ornamental, revegetation, and agroforestry purposes (Turner et al. 1998, Serbesoff-King 2003, Dray 2003, in Pratt et al. 2005b).

Principal source: [Pacific Island Ecosystems at Risk \(PIER\)](#).

Compiler: IUCN/SSC Invasive Species Specialist Group (ISSG)

Review: Dr. Lyn Craven, Principal Research Scientist Australian National Herbarium Australia

Publication date: 2010-10-04

ALIEN RANGE

Global Invasive Species Database (GISD) 2026. Species profile *Melaleuca quinquenervia*. Available from: <https://iucngisd.org/gisd/species.php?sc=45> [Accessed 06 March 2026]

- | | |
|--------------------------------------|------------------------------|
| [1] ANTIGUA AND BARBUDA | [5] BAHAMAS |
| [1] BARBADOS | [1] BENIN |
| [1] BERMUDA | [1] BRAZIL |
| [1] CHINA | [2] COSTA RICA |
| [2] CUBA | [1] DOMINICAN REPUBLIC |
| [1] EGYPT | [1] FIJI |
| [1] FRANCE | [1] FRENCH GUIANA |
| [2] FRENCH POLYNESIA | [1] GRENADA |
| [1] GUADELOUPE | [1] GUAM |
| [1] GUYANA | [1] HAITI |
| [1] HONDURAS | [1] HONG KONG |
| [1] INDIA | [1] ITALY |
| [1] JAMAICA | [1] MADAGASCAR |
| [1] MARTINIQUE | [2] MEXICO |
| [2] MICRONESIA, FEDERATED STATES OF | [1] MONTSERRAT |
| [1] NEW ZEALAND | [1] NICARAGUA |
| [1] PALAU | [1] PANAMA |
| [1] PHILIPPINES | [4] PUERTO RICO |
| [1] REUNION | [1] SAINT LUCIA |
| [1] SAINT VINCENT AND THE GRENADINES | [1] SENEGAL |
| [1] SURINAME | [1] TAIWAN |
| [1] TRINIDAD AND TOBAGO | [1] TURKS AND CAICOS ISLANDS |
| [1] UGANDA | [51] UNITED STATES |
| [2] VIRGIN ISLANDS, U.S. | [1] WEST INDIES |

BIBLIOGRAPHY

84 references found for *Melaleuca quinquenervia*

Management information

Buckingham, Gary R. 2001. Quarantine host range studies with *Lophyrotoma zonalis*, an Australian sawfly of interest for biological control of melaleuca, *Melaleuca quinquenervia*, in Florida, *BioControl (Dordrecht)* 46(3): pp. 363-386.

Burrows, D.W. and Balciunas, J.K. 1997. Biology, distribution and host-range of the sawfly, *Lophyrotoma zonalis* (Hym. Pergidae), a potential biological control agent for the paperbark tree, *Melaleuca quinquenervia*, *Entomophaga* 42(3): pp. 299-313.

Center, T.D., P.D. Pratt, P.W. Tipping, M.B. Rayamajhi, T.K. Van, S.A. Wineriter, F.A. Dray, Jr. & M. Purcell. 2006. Field colonization, population growth, and dispersal of *Boreioglycaspis melaleucae* Moore, a biological control agent of the invasive tree *Melaleuca quinquenervia* (Cav.) Blake, *Biological Control* 39(3): pp. 363-374.

Center, T.D., Pratt, P.D., Tipping, P.W., Rayamajhi, M.B., Van, T.K., Wineriter, S.A. & Dray Jr., F.A. 2007. Initial impacts and field validation of host range for *Boreioglycaspis melaleucae* Moore (Hemiptera: Psyllidae), a biological control agent of the invasive tree *Melaleuca quinquenervia* (Cav.) Blake (Myrtales: Myrtaceae: Leptospermoideae), *Environmental Entomology Volume* 36(3): pp. 569-576.

[Cofrancesco, A.F., Jr. ; Wooten, J.W. ; Jones, H.L. 1995. Evaluation of Mechanical and Chemical Methods for Control of *Melaleuca quinquenervia* in Southern Florida. Wetlands Research Programs. Final Report. Army Engineer Waterways Experiment Station Vicksburg MS.](#)

Summary: Available from: <http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA303356&Location=U2&doc=GetTRDoc.pdf> [Accessed 20 September 2009]

Costello, S.L., Pratt, P.D., Rayamajhi, M.B. & Center, T.D. 2003. Arthropods associated with above-ground portions of the invasive tree, *Melaleuca quinquenervia*, in south Florida, USA, *Florida Entomologist Volume* 86(3): pp. 300-322.

Daehler, C.C; Denslow, J.S; Ansari, S and Huang-Chi, K., 2004. A Risk-Assessment System for Screening Out Invasive Pest Plants from Hawaii and Other Pacific Islands. *Conservation Biology* Volume 18 Issue 2 Page 360.

Summary: A study on the use of a screening system to assess proposed plant introductions to Hawaii or other Pacific Islands and to identify high-risk species used in horticulture and forestry which would greatly reduce future pest-plant problems and allow entry of most nonpests.

Davies, K.A, Makinson, J., Purcell, M.F. 2001. Observations on the development and parasitoids of Fergusonina/Fergusobia galls on *Melaleuca quinquenervia* (Myrtaceae) in Australia, *Transactions of the Royal Society of South Australia* 125(1): pp. 45-50.

Summary: The gall-forming Fergusonina/Fergusobia association is being considered as a potential biocontrol agent of *Melaleuca quinquenervia* in Florida, where it has become a serious weed. This paper reports observations on the development of Fergusonina/Fergusobia galls on *M. quinquenervia* in coastal and sub-coastal south-eastern Queensland and northern New South Wales. The morphology of the gall and the relationship between gall size and numbers of developing cavities and insects are described. Nematodes were found in cavities containing first and second or early third stage fly larvae. Eight species of hymenopteran parasitoids were reared from galls.

[Early Detection and Distribution Mapping System \(EDDMaps\). 2009. *Melaleuca quinquenervia* \(Cav.\) Blake](#)

Summary: Available from: <http://www.eddmaps.org/distribution/usstate.cfm?sub=2783> [Accessed 20 September 2009]

Franks, S.J., Kral, A.M. & Pratt, P.D. 2006. Herbivory by introduced insects reduces growth and survival of *Melaleuca quinquenervia* seedlings *Environmental Entomology* 35(2): pp. 366-372.



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Melaleuca quinquenervia*

- Fuller, D. O. 2005. Remote detection of invasive *Melaleuca* trees (*Melaleuca quinquenervia*) in South Florida with multispectral IKONOS imagery, *International Journal of Remote Sensing* 26(5): pp. 1057 - 1063.
- [Gioeli, T.K. & A.S. Neal. 2004. Biological Control of in Florida s Natural and Man-made Landscapes. Proc. Fla. State Hort. Soc. 117: 254-255.](#)
- Summary:** Available from: <http://www.fshs.org/Proceedings/Password%20Protected/2004%20v.%20117/254-255.pdf> [Accessed 20 September 2009]
- [Hawaii Ecosystems at Risk \(HEAR\). 2008. Images > Myrtaceae > Melaleuca quinquenervia: Paperbark, punk tree](#)
- Summary:** Available from: <http://www.hear.org/starr/plants/images/species/?q=melaleuca+quinquenervia> [Accessed 20 September 2009]
- [IUCN/SSC Invasive Species Specialist Group \(ISSG\)., 2010. A Compilation of Information Sources for Conservation Managers.](#)
- Summary:** This compilation of information sources can be sorted on keywords for example: Baits & Lures, Non Target Species, Eradication, Monitoring, Risk Assessment, Weeds, Herbicides etc. This compilation is at present in Excel format, this will be web-enabled as a searchable database shortly. This version of the database has been developed by the IUCN SSC ISSG as part of an Overseas Territories Environmental Programme funded project XOT603 in partnership with the Cayman Islands Government - Department of Environment. The compilation is a work under progress, the ISSG will manage, maintain and enhance the database with current and newly published information, reports, journal articles etc.
- Kueffer, C. and Mauremootoo, J., 2004. Case Studies on the Status of Invasive Woody Plant Species in the Western Indian Ocean. 3. Mauritius (Islands of Mauritius and Rodrigues). Forest Health & Biosecurity Working Papers FBS/4-3E. Forestry Department, Food and Agriculture Organization of the United Nations, Rome, Italy.
- Laroche, F.B. 1998. Managing melaleuca (*Melaleuca quinquenervia*) in the Everglades, *Weed Technology* 12(4): pp. 726-732.
- Laroche, F.B. 1999. *Melaleuca Management Plan: Ten Years of Successful Melaleuca Management in Florida (1988 - 1998)*. Florida EPPC (Exotic Pest Plant Council)
- [Mack, R. N and W. M. Lonsdale., 2002. Eradicating invasive plants: Hard-won lessons for islands. In Turning the tide: the eradication of invasive species: 311-318. Veitch, C.R. and Clout, M.N.\(eds\). IUCN SSC Invasive Species Specialist Group. IUCN. Gland. Switzerland and Cambridge. UK.](#)
- Summary:** Uses *Clidemia hirta* in Hawaii as an eradication case study. *Clidemia* is in the Melastomataceae and somewhat similar ecologically to *miconia*.
- Eradication case study in Turning the tide: the eradication of invasive species.
- Myers, R.L., H.A. Belles & J.R. Snyder. 2001. Prescribed fire in the management of *Melaleuca quinquenervia* in subtropical Florida. Pages 132-140 in K.E.M. Galley and T.P. Wilson (eds.). Proceedings of the Invasive Species Workshop: the Role of Fire in the Control and Spread of Invasive Species. Fire Conference 2000: the First National Congress on Fire Ecology, Prevention, and Management. Miscellaneous Publication No. 11, Tall Timbers Research Station, Tallahassee, FL.
- [Paciifc Ecosystems at Risk \(PIER\). 2009. Melaleuca quinquenervia \(Cav.\) S.T.Blake. Myrtaceae](#)
- Summary:** Available from: http://www.hear.org/pier/species/melaleuca_quinquenervia.htm [Accessed 20 September 2009]
- [PIER \(Pacific Island Ecosystems at Risk\), 2002. Melaleuca quinquenervia](#)
- Summary:** Ecology, synonyms, common names, distributions (Pacific as well as global), management and impact information. Available from: http://www.hear.org/pier/species/melaleuca_quinquenervia.htm [Accessed 5 February 2003].
- [Plant Conservation Alliance \(PCA\). 2008. Alien Plant Working Group Melaleuca quinquenervia \(Cav.\) Blake Myrtle family \(Myrtaceae\)](#)
- Summary:** Available from: <http://www.nps.gov/plants/alien/fact/mequ1.htm> [Accessed 20 September 2009]
- Pratt, P. D., Slone, D. H., Rayamajhi, M. B., Van, T. K. & Center, T. D. 2003. Geographic distribution and dispersal rate of *Oxyops vitiosa* (Coleoptera: Curculionidae), a biological control agent of the invasive tree *Melaleuca quinquenervia* in south Florida, *Environmental Entomology* 32(2): pp. 397-406.
- Pratt, P. & Ferriter, A. 2001. Plan of Work for The Areawide Management Evaluation of *Melaleuca quinquenervia* (TAME Melaleuca). United States Department of Agriculture.
- Purcell, M. F. & Goolsby, J. A. 2005. Herbivorous insects associated with the paperbark *Melaleuca quinquenervia* and its allies: VI. Pergidae (Hymenoptera), *Australian Entomologist* 32: pp. 37-48.
- Summary:** Surveys were conducted in Australia to find biological control agents for the broad-leaved paperbark tree, *Melaleuca quinquenervia*, a serious pest in Florida, USA. This paper presents collection records and biological information for five sawfly species: *Acanthoperga cameronii* (Westwood), *Perga vollenhovii* Westwood, *Pergagrapta polita* Leach, *Pterygophorus insignis* Kirby and *Lophyrotoma zonalis* (Rohwer); all in family Pergidae. One of these species, *Lophyrotoma zonalis*, was extensively studied as a biological control agent but concerns over its toxicity have delayed release.
- Rayachhetry, M.B., Elliott, M.L., Center, T.D. & Laroche, F. 1999. Field evaluation of a native fungus for control of melaleuca (*Melaleuca quinquenervia*) in southern Florida, *Weed Technology* 13(1): pp. 59-64.
- Rayamajhi, M.B., Van, T.K., Pratt, P.D. & Center, T.D. 2006a. Interactive association between *Puccinia psidii* and *Oxyops vitiosa*, two introduced natural enemies of *Melaleuca quinquenervia* in Florida, *Biological Control* 37(1): pp. 56-67.
- Rayamajhi, Min B., Pratt, Paul D., Center, Ted D., Tipping, Philip W. & Van, Thai K. 2008a. Aboveground biomass of an invasive tree melaleuca (*Melaleuca quinquenervia*) before and after herbivory by adventive and introduced natural enemies: A temporal case study in Florida, *Weed Science* 56(3): pp. 451-456.
- Scheffer, S. J., Giblin-Davis, R. M., Taylor, G. S., Davies, K. A., Purcell, M., Lewis, M. L., Goolsby, J., Center, T. D. 2004. Phylogenetic relationships, species limits, and host specificity of gall-forming Fergusonina flies (Diptera: Fergusoninidae) feeding on *Melaleuca* (Myrtaceae), *Annals of the Entomological Society of America* 97(6): pp.1216-1221.
- Serbesoff-King, Kristina. 2003. *Melaleuca* in Florida: A literature review on the taxonomy, distribution, biology, ecology, economic importance and control measures, *Journal of Aquatic Plant Management* 41: pp. 98-112.
- [SFWMD \(South Florida Water Management District\). Undated. Miami-Dade County Lake Belt.](#)
- Summary:** Available from: http://www.sfwmd.gov/portal/page/portal/pg_grp_sfwmd_watersupply/pg_sfwmd_watersupply_miamidadecolakebelt [Accessed 16 November]
- Silvers, C.S., Pratt, P.D., Ferriter, A.P. & Center, T.D. 2007. TAME melaleuca: A regional approach for suppressing one of Florida s worst weeds, *Journal of Aquatic Plant Management* 45: pp. 1-8.

Stocker, R.K. 1999. Mechanical harvesting of *Melaleuca quinquenervia* in Lake Okeechobee, Florida, *Ecological Engineering* 12(3-4): pp. 373-386.

[TAME \(The Areawide Management Evaluation of *Melaleuca quinquenervia*\). 2007.](#)

Summary: Available from: <http://tame.ifas.ufl.edu/index.shtml> [Accessed 19 November]

Tipping, P.W., Martin, M.R., Nimmo, K.R., Pierce, R.M., Smart, M.D., White, E., Madeira, P.T. & Center, T.D. 2009. Invasion of a West Everglades wetland by *Melaleuca quinquenervia* countered by classical biological control, *Biological Control* 48(1): pp. 73-78.

Tipping, P.W., Martin, M.R., Pratt, P.D., Center, T.D., Rayamajhi, M.B. 2008. Suppression of growth and reproduction of an exotic invasive tree by two introduced insects, *Biological Control* 44(2): 235-241.

Turner, C.E., Center, T.D., Burrows, D.W. & Buckingham, G.R. 1998. Ecology and management of *Melaleuca quinquenervia*, an invader of wetlands in Florida, USA, *Wetlands Ecology and Management* 5(3): pp. 165-178.

[Varnham, K. 2006. Non-native species in UK Overseas Territories: a review. JNCC Report 372. Peterborough: United Kingdom.](#)

Summary: This database compiles information on alien species from British Overseas Territories.

Available from: <http://www.jncc.gov.uk/page-3660> [Accessed 10 November 2009]

Watt, M.S., Kriticos, D.J. & Manning, L.K. 2009. The current and future potential distribution of *Melaleuca quinquenervia*, *Weed Research Volume* 49(4): pp. 381-390.

Wheeler, G.S. & Ordung, K.M. 2006. Lack of an induced response following fire and herbivory of two chemotypes of *Melaleuca quinquenervia* and its effect on two biological control agents, *Biological Control* 39(2): pp. 154-161.

Wineriter, S.A., G.R. Buckingham & F.J. Howard. 2003. Host range of *Boreioglycaspis melaleucae* Moore (Hemiptera: Psyllidae), a potential biocontrol agent of *Melaleuca quinquenervia* (Cav.) S.T. Blake (Myrtaceae), under quarantine, *Biological Control* 27(3): pp. 273-292.

[Wineriter, S.A. & S.E. Halbert. 2002. *Boreioglycaspis melaleucae* \(Hemiptera: Psyllidae\), an Introduced Biocontrol Agent of *Melaleuca quinquenervia* \(Myrtaceae\) in Florida. *Entomology Circular No. 410*. Fl. Dept. Agriculture & Cons. Serv. Division of Plant Industry](#)

Summary: Available from: <http://www.doacs.state.fl.us/pi/enpp/ento/entcirc/ent410.pdf> [Accessed 20 September 2009]

General information

[American Bird Conservancy. 2007. Threats to Birds - *Melaleuca* \(*Melaleuca quinquenervia*\).](#)

Summary: Available from: <http://www.abcbirds.org/conservationissues/threats/invasives/melaleuca.html> [Accessed 20 September 2009]

Bodley, M.J., Ferriter, A.P. & Thayer, D.D. 1994. The biology, distribution, and ecological consequences of *Melaleuca quinquenervia* in the Everglades, *Everglades: the ecosystem and its restoration*: pp. 341-355.

Summary: *Melaleuca* primarily infests the Florida peninsula south of Lake Okeechobee. This area comprises 3 035 000 ha, of which 202 000-607 000 ha) is estimated to be *Melaleuca* infested. Degree of infestation varies from single trees to thousands of trees per acre. A rate of expansion study indicate that the uncontrolled trees could overtake most of the region's remaining natural land within 30 yr. Many public land management agencies are striving to eliminate the plant from their area of responsibility. Herbicidal control is currently the most practical and economically feasible control technique. Biological control may result from the planned winter 1994 release of Australian *Melaleuca*-feeding insects, which may halt the plant's spread by consuming new shoots, seedlings, flowers, and seeds. *Melaleuca* has been declared both a Federal Noxious Weed and a Florida Prohibited Aquatic Plant. -from Authors

[CONABIO. 2008. Sistema de información sobre especies invasoras en México. Especies invasoras - Plantas. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad. Fecha de acceso.](#)

Summary: English:

The species list sheet for the Mexican information system on invasive species currently provides information related to Scientific names, family, group and common names, as well as habitat, status of invasion in Mexico, pathways of introduction and links to other specialised websites. Some of the higher risk species already have a direct link to the alert page. It is important to notice that these lists are constantly being updated, please refer to the main page (<http://www.conabio.gob.mx/invasoras/index.php/Portada>), under the section Novedades for information on updates.

Invasive species - Plants is available from: http://www.conabio.gob.mx/invasoras/index.php/Especies_invasoras_-_Plantas [Accessed 30 July 2008]

Spanish:

La lista de especies del Sistema de información sobre especies invasoras de México cuenta actualmente con información acerca de nombre científico, familia, grupo y nombre común, así como hábitat, estado de la invasión en México, rutas de introducción y ligas a otros sitios especializados. Algunas de las especies de mayor riesgo ya tienen una liga directa a la página de alertas. Es importante resaltar que estas listas se encuentran en constante proceso de actualización, por favor consulte la portada (<http://www.conabio.gob.mx/invasoras/index.php/Portada>), en la sección novedades, para conocer los cambios.

Especies invasoras - Plantas is available from: http://www.conabio.gob.mx/invasoras/index.php/Especies_invasoras_-_Plantas [Accessed 30 July 2008]

[Conservatoire Botanique National De Mascarin \(BOULLET V. coord.\) 2007. *Melaleuca quinquenervia*. - Index de la flore vasculaire de la Réunion \(Trachophytes\) : statuts, menaces et protections. - Version 2007.1](#)

Summary: Base de données sur la flore de la Réunion. De nombreuses informations très utiles.

Available from: <http://flore.cbnm.org/index2.php?page=taxon&num=1dfcb07c683107f038d8c886145d097e> [Accessed March 2008]

Cook, L.G., Morris, D.C., Edwards, R.D. & Crisp, M.D. 2008. Reticulate evolution in the natural range of the invasive wetland tree species *Melaleuca quinquenervia*, *Molecular Phylogenetics & Evolution* 47(2): pp. 506-522.

Craven, L.A. 2006. New Combinations in *Melaleuca* for Australian Species of *Callistemon* (Myrtaceae). *Novon*, 16: 468-475.

Craven, L. In press. *Melaleuca*. In *Flora of Australia*.

Delnatte, pers. comm., 2007

Summary: Personal communication with César Delnatte from the herbarium de Cayenne

Dray, F.A. Jr., Bennett, B.C. & Center, T.D. 2006. Invasion history of *Melaleuca quinquenervia* (Cav.) S.T. Blake in Florida, *Castanea* 71(3): pp. 210-225.

Dray, F.A. Jr., Hale, R.E., Madeira, P.T., Bennett, B.C. & Center, T.D. 2009. Concordance between life history traits, invasion history, and allozyme diversity of the Everglades invader *Melaleuca quinquenervia*, *Aquatic Botany* 90(4): pp. 296-302.



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Melaleuca quinquenervia*

Florence J., Chevillotte H., Ollier C., & Meyer J.-Y. 2007. *Melaleuca quinquenervia*. Base de données botaniques Nadeaud de l'Herbier de la Polynésie française (PAP).

Summary: Base de données sur le flore de Polynésie Française.

Available from: http://www.herbier-tahiti.pf/Selection_Taxonomie.php?id_tax=2742 [Accessed March 2008]

Florida Exotic Pest Plant Council (FLEPPC). Undated. *Melaleuca quinquenervia* (Cav.) Blake

Summary: Available from: http://www.fleppc.org/ID_book/melaleuca%20quinquenervia.pdf [Accessed 18 November 2009]

Geary, T. F. & Woodall, S. L. 1990. *Melaleuca quinquenervia* (Cav.) S. T. Blake *Melaleuca*. In: Burns, Russell M.; Honkala, Barbara H., tech. coords. *Silvics of North America. Vol. 2. Hardwoods. Agric. Handb. 654*. Washington, DC: U.S. Department of Agriculture, Forest Service: 461-465.

Summary: Available from: http://www.na.fs.fed.us/spfo/pubs/silvics_manual/volume_2/melaleuca/quinquenervia.htm [Accessed 3 June, 2010]

Geary, T. Undated. Part II-Species Descriptions: *Melaleuca quinquenervia* (Cav.) S.T. Blake: pp. 567-568.

Summary: Available from: <http://www.rngr.net/publications/ttsm/species/PDF.2004-03-15.2622> [Accessed 3 June, 2010]

Gomes, A.R. Sena and T.T. Kozlowski. 1980. Flooding effects on *Melaleuca quinquenervia*: Responses of *Melaleuca quinquenervia* seedlings to flooding, *Physiol. Plant* 49: pp. 373-377.

Ireland, B.F., Hibbert, D.B., Goldsack, R.J., Doran, J.C. & Brophy, J.J. 2002. Chemical variation in the leaf essential oil of *Melaleuca quinquenervia* (Cav.) S.T. Blake. *Biochemical Systematics and Ecology*, 30(5): 457-470.

ITIS (Integrated Taxonomic Information System). 2005. Online Database *Melaleuca quinquenervia*

Summary: An online database that provides taxonomic information, common names, synonyms and geographical jurisdiction of a species. In addition links are provided to retrieve biological records and collection information from the Global Biodiversity Information Facility (GBIF) Data Portal and bioscience articles from BioOne journals.

Available from:

http://www.cbif.gc.ca/pls/itisca/taxastep?king=every&p_action=containing&taxa=Melaleuca+quinquenervia&p_format=&p_ifx=plgt&p_lang= [Accessed March 2005]

Kaufman, S. and Smouse, P. 2001. Comparing indigenous and introduced populations of *Melaleuca quinquenervia* (Cav.) Blake: response of seedlings to water and pH levels, *Oecologia* 127: pp. 487-494.

Langeland, K.A. and Burks, K. C (Eds) 1998. *Identification and Biology of Non-Native Plants in Florida's Natural Areas*, University of Florida. *Melaleuca quinquenervia*

Summary: Information on plants that pose threats to natural resource areas in Florida.

Available from: http://www.fleppc.org/ID_book/melaleuca%20quinquenervia.pdf [Accessed 30 December 2004]

Lopez-Zamora, I., Comerford, N.B. & Muchovej, R.M. 2004. Root development and competitive ability of the invasive species *Melaleuca quinquenervia* (Cav.) S.T. Blake in the South Florida flatwoods, *Plant and Soil* 263: 239-247.

Mazzotti, F.J., Center, T., Dray, F., Thayer, D. 1997. *Ecological Consequences of Invasion by Melaleuca quinquenervia in South Florida Wetlands: Paradise Damaged, not Lost*. Wildlife Ecology and Conservation Department, University of Florida.

Summary: Available from: <http://edis.ifas.ufl.edu/uw123> [Accessed 21 September 2009]

Munger, Gregory T. 2005. *Melaleuca quinquenervia*. In: *Fire Effects Information System*, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer).

Summary: Available from: <http://www.fs.fed.us/database/feis/plants/tree/melqui/all.html> [Accessed 10 October 2009]

NatureServe. 2009. *NatureServe Explorer: Melaleuca quinquenervia - (Cav.) Blake* *Punktree*

Summary: NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia.

Available from: <http://www.natureserve.org/explorer/servlet/NatureServe?searchName=Melaleuca%20quinquenervia> [Accessed 20 September 2009]

O Hare, N.K. & Dalrymple, G.H. 1997. Wildlife in southern Everglades wetlands invaded by *Melaleuca* (*Melaleuca quinquenervia*), *Bulletin of the Florida Museum of Natural History* 41(1): pp. 1-68.

Summary: In the Everglades region of southeastern Florida, invasion of graminoid/herbaceous wetlands by the invasive, non-native tree *Melaleuca* (*Melaleuca quinquenervia*) results in a closed-canopy forested wetland, with a sparse understory. Intermediate stages in this transformation include a savannah with scattered mature *Melaleuca* trees, and mature dense *Melaleuca* stands surrounded by areas with moderate to low levels of *Melaleuca*. Intermediate levels of *Melaleuca* invasion have not received any attention and were the rationale for our study. Wildlife was surveyed monthly for two years to determine species richness and abundance in wetlands with different *Melaleuca* coverages. Wildlife included all vertebrate classes, as well as selected macro-invertebrates such as crayfish (*Procambarus alleni*) and grass shrimp (*Palaemonetes pugio*). Species richness was highest in areas with moderate *Melaleuca* coverage. Higher species richness is typical of sites with greater vegetative structural diversity, i.e., as in the savannah stage of invasion, as well as areas in an early stage of disturbance. The higher species richness was primarily the result of an increased number of migratory, upland birds. Many of these transient and winter-resident birds occurred at much lower abundances than in native forested habitats such as cypress swamps (*Taxodium distichum*), tropical hardwood hammocks, and pine (*Pinus elliotii* var. *densa*) rocklands. In contrast to the birds, number of species and the abundance of herpetofauna varied little across the *Melaleuca* gradient. There was no shift in species composition from wetland to upland species as the *Melaleuca* coverage increased. The number of fish species was similar across the *Melaleuca* gradient. Unlike the herpetiles, fishes were less abundant in the closed-canopy *Melaleuca* forests, indicating poorer habitat quality. Complex patterns of hydrology and gapping in the forest canopy due to wind storms and fires permitted light penetration and the persistence of productive pockets of aquatic life even within dense stands of *Melaleuca*. The mosaic of areas with low to moderate infestations of *Melaleuca* surrounding mature dense *Melaleuca* stands allowed higher numbers of individuals and species to persist in or seasonally use mature dense *Melaleuca* stands. This interspersed of habitats resulted in stands of *Melaleuca* with ecotonal edges that provided marginal habitat for species characteristic of natural communities. Higher degree of interspersed (more edge) may also mean that the natural areas experience higher exposure to *Melaleuca* seed source, which may result in a faster rate of spread of *Melaleuca*. The results demonstrated that animal populations persisted in areas with disturbed vegetation, as long as critical abiotic factors (in this case hydrology) remained in operation. Areas with moderate levels of *Melaleuca* retained species composition and productivity typical of the natural wetland community. The dominant characteristic of the faunal shifts along the gradient of increasing *Melaleuca* coverage was increased numbers of upland, arboreal, and/or forest species, not the loss of wetland species. Regional permitting and natural resource agencies should recognize that lands with moderate levels of *Melaleuca* may retain significant habitat quality. Restoration of such lands will demonstrate higher levels of success if the method used for *Melaleuca* removal allows for retention of the in situ wildlife community.

Porazinska, D.L., Pratt, P.D. & Giblin-Davis, R.M. 2007. Consequences of *Melaleuca quinquenervia* invasion on soil nematodes in the Florida Everglades, *Journal of Nematology* 39(4): pp. 305-312.

Pratt, P.D., Quevedo, V., Bernier, L., Sustache, J. & Center, T.D. 2005b. Invasions of Puerto Rican wetlands by the Australian tree *Melaleuca quinquenervia*, *Caribbean Journal of Science* 41(1): pp. 42-54.

Summary: Available from: http://www.caribjsci.org/april05/41_42-54.pdf [Accessed 3 June, 2010]

Pratt, P.D., Rayamajhi, M.B., Silvers, C.S. & Ferriter, A.P. 2007. Naturalization and biomass allocation of the invasive tree *Melaleuca quinquenervia* in wetlands of the Bahamas, *Journal of Aquatic Plant Management* 45: pp. 8-16.

Pratt, P.D., Rayamajhi, M. B., Van, T.K., Center, T.D. 2004. Modeling the influence of resource availability on population densities of *Oxyops vitiosa* (Coleoptera: Curculionidae), a biological control agent of the invasive tree *Melaleuca quinquenervia*. *Biocontrol Science and Technology*. 14(1): 51-61.

Summary: Available from: <http://www.informaworld.com/smpp/content~db=all~content=a713616659> [Accessed 3 June, 2010]

Pratt, P.D., Rayamajhi, M.B., Van, T.K., Center, T.D. & Tipping, P.W. 2005a. Herbivory alters resource allocation and compensation in the invasive tree *Melaleuca quinquenervia*, *Ecological Entomology* 30(3): pp. 316-326.

Ramanoelina, P.A.R., Bianchini, J.P. & Gaydou, E.M. 2008. Main industrial niaouli (*Melaleuca quinquenervia*) oil chemotype productions from Madagascar. *Journal of Essential Oil Research*, 20: 261-266.

Summary: Available from: http://www.ism2.univ-cezanne.fr/fichiers_pdf/807.pdf

Rayamajhi, M.B., Pratt, P.D., Center, T.D., Tipping, P.W. & Van, T.K. 2008b. Decline in exotic tree density facilitates increased plant diversity: the experience from *Melaleuca quinquenervia* invaded wetlands, *Wetlands Ecology and Management*: pp. 1-13.

Rayamajhi, M.B., Purcell, M.F., Van, T.K., Center, T.D., Pratt, P.D. & Buckingham, G.R. 2002. 8. Australian Paperbark Tree (*Melaleuca*). In *Biological control of invasive plants in the Eastern United States*. R. G. V. Driesche, B. Blossey, M. S. Hoddle, S. Lyon, and R. Reardon (Eds). Pp. 117-130. Morgantown, West Virginia: Forest Health Technology Enterprise Team.

Summary: Available from: <http://www.invasiveplants.net/biologicalcontrol/pdf/8AustralianPaperbarkTree.pdf> [Accessed 3 June, 2010]

Rayamajhi, M.B., Van, T.K., Pratt, P.D. & Center, T.D. 2006b. Temporal and structural effects of stands on litter production in *Melaleuca quinquenervia* dominated wetlands of south Florida, *Wetlands Ecology and Management Volume* 14(4): pp. 303-316.

Rayamajhi, Min B., Van, Thai K., Pratt, Paul D., Center, Ted D. & Tipping, Phillip W. 2007. *Melaleuca quinquenervia* dominated forests in Florida: analyses of natural-enemy impacts on stand dynamics, *Plant Ecology* 192(1): pp. 119-132.

Scher, Julia., 2009. Federal Noxious Weed Disseminules of the US. *Melaleuca quinquenervia* (Cav.) S.T. Blake Family: Myrtaceae

Summary: Available from: <http://keys.lucidcentral.org/keys/FNW/FNW%20seeds/html/fact%20sheets/Melaleuca%20quinquenervia.htm> [Accessed 20 September 2009]

USDA-ARS. 2009. Taxon: *Melaleuca quinquenervia* (Cav.) S. T. Blake National Genetic Resources Program. Germplasm Resources Information Network - (GRIN) [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland.

Summary: Available from: <http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?105723> [Accessed 20 September 2009]

USDA, NRCS. 2009. The PLANTS Database *Melaleuca quinquenervia* (Cav.) S.F. Blake punktree

Summary: Available from: <http://plants.usda.gov/java/profile?symbol=mequ> [Accessed 20 September 2009]

Van, Thai. K., Min, B. Rayamajhi & Ted, D. Center. 2005. Seed Longevity of *Melaleuca quinquenervia*: A Burial Experiment in South Florida, *J. Aquat. Plant Manage* 43: pp. 39-42.

Van, T.K., Rayachhetry, M. B. & Center, T. D., 2000. Estimating above-ground biomass of *Melaleuca quinquenervia* in Florida, USA, *Journal of Aquatic Plant Management* 38: pp. 62-67.

Summary: Available from: <http://aquacomm.fcla.edu/1839/1/v38p62.pdf> [Accessed 20 September 2009]



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Melaleuca quinquenervia*

- Wagner, W. L., Derral, R. H. and Sohmer, S. H. 1990. Manual of the flowering plants of Hawaii i: 373-374.
- Wilson, P.G., O'Brien, M.M., Heslewood, M.M. & Quinn, C.J. 2005. Relationships within Myrtaceae sensu lato based on a matK phylogeny. *Plant Systematics and Evolution*, 251: 3-19.
- Woodall, S.L. 1981. Site requirements for melaleuca seedling establishment. In: Geiger, R. K., comp. *Proceedings of melaleuca symposium*; 1980 September 23-24; [Location unknown]. Tallahassee, FL: Florida Division of Forestry: 9-15.