

Acacia farnesiana  [简体中文](#) [正體中文](#)

System: Terrestrial

| Kingdom | Phylum | Class | Order | Family |
|---------|---------------|---------------|---------|----------|
| Plantae | Magnoliophyta | Magnoliopsida | Fabales | Fabaceae |

Common name cashia (English, Puerto Rico), Ellington curse (English), vaivai vakavotona (English, Fiji), aromo (Spanish, Spain), ban baburi (English, Fiji), mimosa (Spanish, Spain), huisache (English, Mexico), esponjeira (Portuguese, Portugal), espino blanco (English, El Salvador), carambuco (Spanish, Spain), espino ruco (English, El Salvador), rayo (English, Puerto Rico), opoponax (English, southern United States), Westindische akazie (German), mimosa bush (English, Australia), needle bush (English, Australia), debena (English, Nauru), tekaibakoa (English, Kiribati), oki (English, Fiji), kolu (English, Hawaii), popinac (English, Guam), aroma (English, Guam), kandaroma (English, Commonwealth of the Northern Mariana Islands), popinac (English, Commonwealth of the Northern Mariana Islands), titima (English, Cook Islands), acacia jaune (French), klu (English), sweet acacia (English), huisache dulce (Spanish), Small's acacia (English), klu bush (English), cassie (French), esponja (Portuguese, Brazil)

Synonym *Mimosa farnesiana*, L.
Mimosa acicularis, Poir.
Acacia minuta, (M.Jones) Beauchamp subsp. *densiflora* (Alex. ex Small) Beauchamp
Mimosa indica, Pers.
Acacia indica, (Pers.) Desv.
Mimosa farnesiana, L.
Vachellia densiflora, Alex. ex Small
Acacia smallii, Isely
Mimosa acicularis, (Humb. & Bonpl. ex Willd.) Poir.
Mimosa edulis, (Humb. & Bonpl. ex Willd.) Poir.
Vachellia farnesiana, (L.) Wight & Arn. forma *typica* Speg.
Vachellia farnesiana, (L.) Wight & Arn. var. *typica* Speg.
Vachellia farnesiana, (L.) Wight & Arn.
Mimosa pedunculata, (Willd.) Poir.
Acacia farnesiana, (L.) Willd. var. *pedunculata* (Willd.) Kuntze
Acacia pedunculata, Willd.
Acacia ferox, M. Martens & Galeotti
Farnesia odora, Gasp.
Acacia acicularis, Humb. & Bonpl. ex Willd.
Acacia lenticellata, F. Muell.
Acacia densiflora, (Alex. ex Small) Cory
Acacia edulis, Humb. & Bonpl. ex Willd.

Similar species *Acacia nilotica*, *Prosopis*

Summary Probably a native of tropical America, *Acacia farnesiana* was introduced to many tropical countries for its bark, gum, seed and wood. It is often planted as an ornamental or to check erosion, and is also used in the perfume industry because of its scented flowers. This thorny, deciduous shrub grows to 4m in height forming impenetrable thickets or sometimes a more open cover and prefers dry habitats between sea level and 1000 m. In Australia it occurs along watercourses on rangeland and farmland limiting access to water. It has also become an invasive species in Fiji, French Polynesia, New Caledonia, Solomon Islands, and Vanuatu.



[view this species on IUCN Red List](#)

Species Description

"This thorny, deciduous shrub grows to 4 metres in height, sometimes forming impenetrable thickets, although in most areas it forms a more open cover" (Smith, 1985. In PIER, 2002). "Erect much-branched shrub; leaves with 4-8 pairs of pinnae, pinnae with 10-12 pairs of small asymmetric leaflets ; stipular thorns straight and slender; flowers in pedunculate axillary heads, 1-3 heads together, subglobose; flowers yellow, fragrant; heads about 1-1.5cm across; stamens numerous; pods dark brown or black, up to 8cm long, to 12mm broad, plump, often slightly curved; pulp within sweetish; seeds compressed, elliptic, brown" (Stone, 1970, in PIER, 2003) (differences according to Paiva, 1999).

Notes

In Hawaii *Acacia farnesiana* was formerly cultivated for an attempted perfume industry. It is now naturalised and common, sometimes becoming a pest in dry, open, disturbed areas, 2-400m (Wagner *et al.*, 1999. In PIER, 2002). Although the aerial portions of the plant may be killed by fire, it soon regenerates from basal shoots, (Smith, 1985. In PIER, 2002). In Spain *Acacia farnesiana* is naturalised in rocky, poor soils (Paiva, 1999)

Uses

A tree of economic importance in South and East Africa, Rhodesia, India and the Rio Grande do Sul area of South America (Duke, 1983). The bark and the pods are a source of tannin and are used for tanning and dyeing leather (University of Connecticut, 2003). The flowers provide a fragrant essential oil which is used in the perfume industry as a violet scent substitute (Le Hou'erou, 2002). A gummy substance obtained from the pods is used in Java as cement for broken crockery. Other parts of the plant are used as an ingredient in the Ivory Coast for arrow poison (University of Connecticut, 2003). Trees add nitrogen and organic material, which improve the soil and are sometimes used for erosion control on poor sloping soils unsuitable for agriculture (Duke, 1983). Products are often used in folk medicine as styptics or astringents (Duke, 1983). In India and some African countries the pods are used as substitute for tamarind *Tamarindus indica* L. (Paiva, 1999). Cultivated in S.W. Europe for ornamental purposes and for perfumery industry (Tutin *et al.*, 1992).

Habitat Description

Acacia farnesiana thrives in dry localities and on loamy or sandy soils where it may serve as a sand binder (ranging from warm temperate dry through tropical desert to moist forest life zones, *Acacia farnesiana* is reported to tolerate annual precipitation of 6.4 - 40.3 dm (a mean of 20 cases is 14.0 dm), annual mean temperature of 14.7-27.8°C (mean of 20 cases = 24.1°C), and pH of 5.0-8.0 (mean of 15 cases = 6.8) (Duke, 1983). Dry habitats between sea level and 1000m (PIER, 2002). Smith (1985 In PIER, 2002) reports that although the aerial portions may be killed by fire, this plant soon regenerates from basal shoots.

Reproduction

It is a prolific seed producer (Scifres, 1974). The seeds readily germinate after soil disturbance and plants grow rapidly (Mutz *et al.*, 1978, Land Protection, 2001). Seeds are dispersed by ungulates which eat the pods (PIER, 2002).

Nutrition

Acacia farnesiana can fix atmospheric nitrogen through symbiotic relation with *Rhizobium* allowing growth in nitrogen poor soils. Thrives on poor, dry soils but favours deeper, moister, more fertile soils (Duke, 1983).



General Impacts

Acacia farnesiana can spread readily and grow quickly, (Land Protection, 2001). As it often forms thorny thickets along some watercourses in Australia, it can be a considerable nuisance during mustering and can also hinder access to water (Land Protection, 2001). While access is less of a problem in areas where cattle graze on the mimosa, they readily eat the nutritious seed pods assisting its spread.

Management Info

Preventative measures: A [Risk Assessment of *Acacia farnesiana*](#) for Hawai'i and other Pacific islands was prepared by Dr. Curtis Daehler (UH Botany) with funding from the Kaulunani Urban Forestry Program and US Forest Service. The alien plant screening system is derived from Pheloung *et al.* (1999) with minor modifications for use in Pacific islands (Daehler *et al.* 2004). The result is a score of 14 and a recommendation of: "Likely to cause significant ecological or economic harm in Hawai'i and on other Pacific Islands as undetermined by a high WRA score, which is based on published sources describing species biology and behaviour in Hawai'i and/or other parts of the world."

Physical: "Destroyed by cultivation and grubbing" (Swarbrick, 1997. In PIER, 2002). Although the aerial portions may be killed by fire, it soon regenerates from basal shoots" (Smith, 1985). Once established, the seedlings grow rapidly and resprout readily following damage or top removal. *Acacia farnesiana* sprouts may grow to almost half their original total plant height within 5 months after shredding (Powell *et al.*, 1972). Thus, mechanical top removal results in only short-term suppression of *A. farnesiana* (Mutz *et al.* 1978) and gives the species competitive advantage over associated, slower growing woody plants. Almost pure, dense stands of *A. farnesiana* may develop within two to three growing seasons following brush control methods that disturb the soil.

Chemical: Probably susceptible to translocated herbicides, including picloram, metsulfuron-methyl, glyphosate and triclopyr, and possibly 2,4-D applied to the foliage, freshly cut stumps or by stem injection at standard rates.

Also probably susceptible to residual herbicides, including tebuthiuron and hexazinone (Swarbrick, 1997 in PIER, 2002). "Sensitive to foliar applications of triclopyr at 1 lb/acre and metsulfuron at 0.45 oz/a and to basal bark applications of 2,4-D or triclopyr at 2% in diesel. Drizzle applications were not effective in foliar and basal bark trials at Kihei, Maui, but these trials were confounded by a severe drought" (Motooka *et al.* 2002. In PIER, 2002).

Pathway

In Hawaii, formerly cultivated for an attempted perfume industry (Wagner *et al.*, 1999. In PIER, 2002)

Principal source: [Pacific Islands Ecosystems at Risk, \(PIER, 2002\) *Acacia farnesiana*](#)

Compiler: IUCN/SSC Invasive Species Specialist Group (ISSG)

Review: Dr. Hélia Marchante. Escola Superior Agrária de Coimbra Departamento de Ciências Exactas e Ambiente Sector de Biologia e Ecologia, Bencanta. 3040-316 Coimbra Portugal.

Publication date: 2006-03-23

ALIEN RANGE

| | |
|-----------------------|-------------------------|
| [1] ANGUILLA | [1] ANTIGUA AND BARBUDA |
| [1] ARABIAN PENINSULA | [6] AUSTRALIA |
| [1] BAHAMAS | [1] BARBADOS |
| [1] CAMBODIA | [1] CAYMAN ISLANDS |
| [1] CHINA | [2] COOK ISLANDS |
| [1] CUBA | [1] DOMINICA |
| [1] EAST AFRICA | [1] EASTERN AFRICA |

| | |
|------------------------------|---------------------|
| [2] FIJI | [2] FRANCE |
| [3] FRENCH POLYNESIA | [1] GRENADA |
| [1] GUADELOUPE | [1] GUAM |
| [1] INDIA | [1] INDONESIA |
| [2] ITALY | [1] JAMAICA |
| [2] JAPAN | [1] KENYA |
| [2] KIRIBATI | [1] MALAYSIA |
| [1] MARTINIQUE | [1] MAURITIUS |
| [1] MAYOTTE | [1] MONTSERRAT |
| [1] NAURU | [2] NEW CALEDONIA |
| [1] NIUE | [1] NORTHERN AFRICA |
| [2] NORTHERN MARIANA ISLANDS | [1] PAKISTAN |
| [1] PHILIPPINES | [1] PORTUGAL |
| [1] PUERTO RICO | [1] REUNION |
| [1] SAHEL | [1] SAINT HELENA |
| [1] SEYCHELLES | [1] SOLOMON ISLANDS |
| [1] SOUTH AMERICA | [1] SPAIN |
| [1] TAIWAN | [1] THAILAND |
| [2] UNITED STATES | [1] VANUATU |
| [1] VIET NAM | |

BIBLIOGRAPHY

30 references found for *Acacia farnesiana*

Management information

Arévalo, José Ramón; Leila Afonso; Agustín Naranjo and Marcos Salas, 2010. Invasion of the Gran Canaria ravines ecosystems (Canary Islands) by the exotic species *Acacia farnesiana*. Plant Ecology. Volume 1 / 1948 - Volume 210 / 2010

[Bontrager, O.E., Scifres, C.J. and Drawe, D.L. 1979. Huisache Control by Power Grubbing. Journal of Range Management, 32\(3\): 185-188.](#)

Summary: Management information by Power Grubbing.

Available from: <http://jrm.library.arizona.edu/data/1979/323/6bont.pdf> [Accessed 10 October 2002].

Daehler, C.C; Denslow, J.S; Ansari, S and Huang-Chi, K., 2004. A Risk-Assessment System for Screening Out Invasive Pest Plants from Hawaii and Other Pacific Islands. Conservation Biology Volume 18 Issue 2 Page 360.

Summary: A study on the use of a screening system to assess proposed plant introductions to Hawaii or other Pacific Islands and to identify high-risk species used in horticulture and forestry which would greatly reduce future pest-plant problems and allow entry of most nonpests.

[European and Mediterranean Plant Protection Organization \(EPPO\), 2006. Guidelines for the management of invasive alien plants or potentially invasive alien plants which are intended for import or have been intentionally imported. EPPO Bulletin 36 \(3\), 417-418.](#)

Land Protection, 2001. pest series Mimosa bush (*Acacia farnesiana*). The State of Queensland, Department of Natural Resources and Mines, June 2001.

[Le Hourou, Acacia farnesiana \(L.\) Willd. Grassland Index. UN FAO.](#)

Summary: Drawings, description, common names, habitat, soil, propagation, distribution, crop management, products and uses, links, references, photos.

Available from: <http://www.fao.org/ag/AGP/AGPC/doc/Gbase/DATA/Pf000113.HTM> [Accessed on 10 October 2002].

Motooka, P., L. Castro, D. Nelson, G. Nagai and L. Ching . 2002. Weeds of pastures and natural areas of Hawaii and their management. In press.

[Rasmussen, G.A., Scifres, C.J. and Drawe, D.L. 1983. Huisache Growth, Burning Quality, and Use Browse Following. Journal of Range Management 36\(3\): 337:342.](#)

Summary: Huisache Growth, Burning Quality, and Use Browse Following.

Available from: <http://jrm.library.arizona.edu/data/1983/363/15rasm.pdf> [Accessed 10 October 2002].

[Varnham, K. 2006. Non-native species in UK Overseas Territories: a review. JNCC Report 372. Peterborough: United Kingdom.](#)

Summary: This database compiles information on alien species from British Overseas Territories.

Available from: <http://www.jncc.gov.uk/page-3660> [Accessed 10 November 2009]

General information

Barthelat, F. 2005. Note sur les espèces exotiques envahissantes Mayotte. Direction de l'Agriculture et de la Forêt. 30p

Summary: Tableau synthétique des plantes exotiques de Mayotte classées en fonction de leur niveau d'envahissement.

[Chihuahuan Desert website, 2002. Sweet Acacia \(Acacia farnesiana\).](#)

Summary: Includes common names and scientific names, plus distribution.

Available from: <http://nasa.utep.edu/chih/gardens/plants/acaciafarn.htm> [Accessed 27 November 2002].

[Conservatoire Botanique National De Mascarin \(BOULLET V. coord.\) 2007. - Acacia farnesiana Index de la flore vasculaire de la Réunion \(Trachophytes\) : statuts, menaces et protections. - Version 2007.1](#)

Summary: Base de données sur la flore de la Réunion. De nombreuses informations très utiles.

Available from: <http://flore.cbnm.org/index2.php?page=taxon&num=6c8349cc7260ae62e3b1396831a8398f> [Accessed 28 March 2008]

Deyrup, M., Davis, L. and Cover, S. 2000. Exotic ants in Florida. Transactions of the American Entomological Society 126: 293-326.

Summary: More than 50 species of exotic ants have established breeding populations in Florida. This is the largest exotic ant fauna of any U.S. state. An annotated list of species (including distribution, origin, and pest status) includes 4 new records for the U.S.

Duke, James A. 1983. Handbook of Energy Crops. Unpublished. *Acacia mearnsii*. Center for New Crops & Plant Products, Purdue University.

Summary: Uses, Folk Medicine, Chemistry, Description, Germplasm, Distribution, Ecology, Cultivation, Harvesting, Yields and Economics, Energy, Biotic Factors, References.

Available from: http://www.hort.purdue.edu/newcrop/duke_energy/Acacia_mearnsii.html [Accessed 10 October 2002].

Fernald, H. T. 1947. The little fire ant as a house pest. J. Econ. Entomol. 40: 428.

Florence J. Chevillotte H. Ollier C. & Meyer J.-Y. 2007. *Acacia farnesiana* Base de données botaniques Nadeaud de l'Herbier de la Polynésie française (PAP).

Summary: Base de données sur la flore de Polynésie Française.

Available from: http://www.herbier-tahiti.pf/Selection_Taxonomie.php?id_tax=2445 [Accessed 26 March 2008]

Franco, J.A. 1943. Dendrologia Florestal. Lisboa.

Summary: Dendrology book of species present in Portugal.

Gargominy, O., Bouchet, P., Pascal, M., Jaffre, T. and Tourneau, J. C. 1996. Consequences des introductions d'espèces animales et végétales sur la biodiversité en Nouvelle-Calédonie. Rev. Ecol. (Terre Vie) 51: 375-401.

Summary: Consequences to the biodiversity of New Caledonia of the introduction of plant and animal species.

ITIS (Integrated Taxonomic Information System). 2004. Online Database *Acacia farnesiana*

Summary: An online database that provides taxonomic information, common names, synonyms and geographical jurisdiction of a species. In addition links are provided to retrieve biological records and collection information from the Global Biodiversity Information Facility (GBIF) Data Portal and bioscience articles from BioOne journals.

Available from: http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=26427 [Accessed December 31 2004]

Meyer, Jean-Yves & Loope, Lloyd & Sheppard, A. & Munzinger, Jérôme & Jaffré, Tanguy. (2006). Les plantes envahissantes et potentiellement envahissantes dans l'archipel néo-calédonien : première évaluation et recommandations de gestion.

Summary: Synthèse sur les plantes exotiques envahissantes et potentiellement envahissantes en Nouvelle-Calédonie. Des recommandations de gestion sont également proposées.

Meyer, J.-Y. 2000. Invasive plants in the Pacific Islands. In: The Invasive Species in the Pacific: A Technical Review and Draft Regional Strategy. Sherley, G. (tech. ed). Published in June 2000 by the South Pacific Regional Environment Programme (SPREP).

Summary: Resource that includes the distribution of invasive species throughout the Pacific Islands.

Meyer, J.-Y. 2007. Conservation des forêts naturelles et gestion des aires protégées en Polynésie française. Bois et forêts des tropiques, 291 (1), 25-30.

Mutz, J.L., Scifres, C.J., Drawe, D.L., Box, T.W. and Whitson, R.E. 1978. Range vegetation after mechanical brush treatments on the Coastal Prairie. Texas Agr. Exp. Sta. Bull. 1191.

Paiva, J. 1999. *Acacia*. In Talavera, S. Aedo, C, Castroviejo, S, Romero Zarco, C, Sorez, L, Salgueiro, F.J & Velayos, (ed). Flora Iberica - Plantas Vasculares de la Península Ibérica e Islas Baleares. Vol.VII(I). Leguminosae. Real Jardón Botánico, CSIS. Madrid. ISBN 84-00-06221-3. pp. 11- 25.

Summary: Flora of Iberian Peninsula.

Plant Gene Resources of Canada (PGRC). Research Branch of Agriculture and Agri-Food Canada.

Summary: Available from: <http://pgrc3.agr.ca/cgi-bin/npgs/html/taxon.pl?844> [Accessed 14 December, 2003].

Scifres, C.J. 1974. Salient aspects of huisache seed germination. Southw. Natur. 18:383-392.

Space, J. C. 2000. Pacific Island Ecosystems at Risk (PIER). *Acacia farnesiana*

Summary: Fact Sheet.

Tassin, J., Rivière, J.N., Cazanove, M., Bruzseses, E. 2006. Ranking of invasive woody plant species for management on Réunion Island. Weed research 46, 388-403

Summary: L'inventaire de 318 espèces de plantes ligneuses introduites à la Réunion, permet d'en identifier 132 comme naturalisées dans les écosystèmes naturels. 26 de ces espèces choisies parmi les plus envahissantes ont été classées en fonction de leur impact biologique sur les écosystèmes indigènes.

University of Connecticut. *Acacia farnesiana* (L.) Willd. Ecology & Evolutionary Biology Conservatory.

Summary: Common Names, Habitat, Description, Uses, Classification.

Available from: http://florawww.eeb.uconn.edu/acc_num/198500174.html [Accessed on 07 January 2003].

USDA-NRCS (Natural Resource Conservation Service). 2005. *Acacia farnesiana*. The PLANTS Database Version 3.5 [Online Database] National Plant Data Center, Baton Rouge, LA.

Summary: Available from: <http://plants.usda.gov/java/nameSearch?mode=Scientific+Name&keywordquery=Acacia+farnesiana> [Accessed 12 March 2006].