

Hemigrapsus sanguineus

System: Marine_terrestrial

Kingdom	Phylum	Class	Order	Family
Animalia	Arthropoda	Malacostraca	Decapoda	Varunidae

Common name Asian shore crab (English), Japanese shore crab (English)

Synonym

Similar species

Summary *Hemigrapsus sanguineus* is commonly called the Asian shore crab and is native to the Asia-Pacific region. It has a very broad diet and, in its introduced range, has the potential to affect populations of native species, such as crabs, fish and shellfish by disrupting the food web. It also occupies habitats very similar to native mud crabs. *Hemigrapsus sanguineus* may compete with larger species, like the blue crab, rock crab and the non-native green crab.



[view this species on IUCN Red List](#)

Species Description

Hemigrapsus sanguineus has several distinguishing characteristics. It has a somewhat square carapace which is usually patterned and dark in colour, ranging from brownish orange to greenish black. Each side of the carapace has three distinct anterolateral teeth. *H. sanguineus* has a readily observable banding pattern on the walking legs. Male crabs possess a fleshy knob at the base of the dactyl of the cheliped, as well as relatively larger and more robust claws than females (Ledesma and O'Connor, 2001). The abdomen of mature females is wider than that of male crabs (Mcdermott 1998a). Benson (2005) states that, "This species is small with adults ranging from 35mm (1.5 in) to 42mm (1.65 in) in carapace width."

Notes

In their studies, Jensen *et al.* (2002) found that, "When competing for food against *H. sanguineus*, the non-native green crab (*Carcinus maenas*) was usually the first to find the bait yet was almost invariably dislodged shortly thereafter by *H. sanguineus*. This typically did not involve overt aggression or chelae displays on the part of the *H. sanguineus*, but rather the use of the body and legs as a wedge to displace the *C. maenas* from the food. *H. sanguineus* rarely used their chelae for grasping their opponent or defending the bait; rather, approaching *C. maenas* were fended off with kicks from the walking legs while the chelae continued to be used for feeding."

Lifecycle Stages

Benson (2005) reports that in its introduced range, "*Hemigrapsus sanguineus* is highly reproductive with a breeding season from May to September, twice the length of native crabs. The females are capable of producing 50,000 eggs per clutch with 3-4 clutches per breeding season. The larvae are suspended in the water for approximately one month before developing into juvenile crabs." Ledesma and O'Connor (2001) report that, "Life history studies suggest that *H. sanguineus* has a high reproductive potential, facilitating a rapid rate of dispersal and colonization of new areas (Fukui 1988; Mcdermott 1991, 1998a)."

Habitat Description

Hemigrapsus sanguineus inhabits any shallow hard-bottom intertidal or sometimes subtidal habitat, with the highest densities occurring at the middle and lower intertidal elevations. It can live on artificial structures and on mussel beds and oyster reefs, and is usually found under the shelter of rocks, shells and other debris on tidal flats. It will search along the fringe of tidal flats as the tide recedes for unused burrows to occupy. Lohrer *et al.* (2000) found that the availability of shelter sites influenced the density and distribution of *H. sanguineus*. *H. sanguineus* can tolerate a wide range of salinities and temperature, as well as damp conditions in the upper intertidal regions (Benson, 2005). It coexists with mud crabs in the family Xanthidae, juvenile green crabs (*Carcinus maenas*), and occasionally the rock crab, *Cancer irroratus* (Benson, 2005; Brousseau *et al.* 2003; Ledesma and O'Connor, 2001).

The primary habitat of *H. sanguineus* in Japan is crevices among boulders on rocky intertidal shores (Fukui, 1988; in Gerard *et al.* 1999). In the Western Pacific the distribution of *H. sanguineus* ranges from ~20 to 50 N latitude including Hong Kong and Taiwan, the Chinese and Korean coasts, and all of Japan (Mcdermott, 1998).

Reproduction

Gerard *et al.* (1999) states that "The reproductive output of this species is impressive: mature females may have two or more broods annually, with a mean clutch size of 15,000 and a maximum of at least 50,000 eggs per crab (Fukui 1988; McDermott 1998b). Planktonic larval stages, which last for about a month under optimal temperature and salinity conditions (Epifanio *et al.* 1998), provide a mechanism for dispersal. Growth and maturation are rapid: newly settled juveniles have a mean carapace width (CW) around 2mm and reach 20mm in about two years (Fukui, 1988). The crabs become reproductively mature at this age and, although growth is slower in mature crabs, they can reach a maximum CW of 40mm, corresponding to a maximum lifespan of around eight years."

Ledesma and O'Connor (2001) states that, "Mcdermott (1998a) suggested that the length of the reproductive period of *H. sanguineus* is related to latitude and therefore water temperature. In southern parts of Japan, the breeding season is 8 months long (Fukui 1988), whereas in northern Japan, it lasts 3 months (Takahashi *et al.* 1985). Whereas breeding occurred through September at Gooseberry in Buzzards Bay, ovigerous females were found only until early August in Sandwich in Cape Cod Bay."

Nutrition

Gerard *et al.* (1999) states that "Like many grapsid crabs, *Hemigrapsus sanguineus* is an opportunistic omnivore, consuming a wide variety of plant and animal foods in the field and in captivity (Fukui 1988; McDermott 1991, 1998b; Takahashi and Matsuura 1994; Lohrer and Whitlatch 1997)."

General Impacts

Hemigrapsus sanguineus has the potential to cause significant changes in the inshore marine and estuarine communities of southern New England and mid-Atlantic coast, and predation by *H. sanguineus* on several species important to the structure of Atlantic rocky intertidal communities provides the strongest evidence to date that *H. sanguineus* has the potential for significant ecological impact in its introduced range (Gerard *et al.* 1999).

The intertidal distribution of *H. sanguineus* overlaps that of juvenile (non-native) green crabs (*Carcinus maenas*), particularly in the northern part of their range. Furthermore, *H. sanguineus* is dependent on rock cover and anecdotal reports suggest *C. maenas* has become uncommon under rocks in some areas since the arrival of *H. sanguineus* (Jensen *et al.* 2002). It is debatable whether there is evidence that *H. sanguineus* is negatively impacting on native crabs (G. Jensen, pers. comm.). Brousseau *et al.* (2003) found that mud crabs were actually competitively dominant over *H. sanguineus*.

Laboratory studies have shown that *H. sanguineus* readily consumes three species of commercial bivalves: blue mussels *Mytilus edulis*, soft-shell clams *Mya arenaria*, and oysters *Crassostrea virginica* (Brousseau *et al.* 2001; Benson, 2005). The high densities of *H. sanguineus* which occur in the wild, their effectiveness as predators of juvenile bivalves and their large appetites suggest an important role for these predators in restructuring the prey communities in habitats into which they have been introduced (Brousseau *et al.* 2001).



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Hemigrapsus sanguineus*

Management Info

Preventative measures: Ballast water management is being researched to reduce or eradicate new introductions from occurring. Scientists are monitoring changes in native species, tracking the shore crab's spread along the US Atlantic coastline, and conducting experiments to increase their knowledge of basic biology and ecology of this species (Benson, 2005).

Biological: It is possible that species such as tautogs, drums, sheepshead and seagulls prey upon *H. sanguineus* (Benson, 2005; G. Jensen, pers. comm.). Parasites, which help control populations of *H. sanguineus* in its native range, are not present along the US Atlantic coast (Benson, 2005).

Principal source: Brousseau *et al.* 2001. *Laboratory investigations of the effects of predator sex and size on prey selection by the Asian crab, Hemigrapsus sanguineus.*

[Benson, 2005](#). Nonindigenous Species Information Bulletin Asian shore crab, Japanese shore crab, Pacific crab, *Hemigrapsus sanguineus* (De Haan) (Arthropoda: Grapsidae).

Compiler: National Biological Information Infrastructure (NBII) & IUCN/SSC Invasive Species Specialist Group (ISSG)

Review: Dr Greg Jensen

Publication date: 2006-11-16

ALIEN RANGE

[1] CROATIA

[1] MEDITERRANEAN & BLACK SEA

[27] UNITED STATES

[2] FRANCE

[1] NETHERLANDS

BIBLIOGRAPHY

32 references found for *Hemigrapsus sanguineus*

Management information

[Benson, A. 2005. Nonindigenous Species Information Bulletin: Asian shore crab, Japanese shore crab, Pacific crab, Hemigrapsus sanguineus \(De Haan\) \(Arthropoda: Grapsidae\). USGS-FISC: Center for Aquatic Resource Studies.](#)

Summary: Available from: http://cars.er.usgs.gov/Nonindigenous_Species/Asian_shore_crab/asian_shore_crab.html [Accessed 13 May 2005]
[Centre for Environment, Fisheries & Aquaculture Science \(CEFAS\), 2008. Decision support tools-Identifying potentially invasive non-native marine and freshwater species: fish, invertebrates, amphibians.](#)

Summary: The electronic tool kits made available on the Cefas page for free download are Crown Copyright (2007-2008). As such, these are freeware and may be freely distributed provided this notice is retained. No warranty, expressed or implied, is made and users should satisfy themselves as to the applicability of the results in any given circumstance. Toolkits available include 1) FISK- Freshwater Fish Invasiveness Scoring Kit (English and Spanish language version); 2) MFISK- Marine Fish Invasiveness Scoring Kit; 3) MI-ISK- Marine invertebrate Invasiveness Scoring Kit; 4) FI-ISK- Freshwater Invertebrate Invasiveness Scoring Kit and AmphISK- Amphibian Invasiveness Scoring Kit. These tool kits were developed by Cefas, with new VisualBasic and computational programming by Lorenzo Vilizzi, David Cooper, Andy South and Gordon H. Copp, based on VisualBasic code in the original Weed Risk Assessment (WRA) tool kit of P.C. Pheloung, P.A. Williams & S.R. Halloy (1999).

The decision support tools are available from:

<http://cefas.defra.gov.uk/our-science/ecosystems-and-biodiversity/non-native-species/decision-support-tools.aspx> [Accessed 13 October 2011]

[The guidance document](http://www.cefas.co.uk/media/118009/fisk_guide_v2.pdf) is available from http://www.cefas.co.uk/media/118009/fisk_guide_v2.pdf [Accessed 13 January 2009].

[Hayes, K., Sliwa, C., Migus, S., McEnulty, F., Dunstan, P. 2005. National priority pests: Part II Ranking of Australian marine pests. An independent report undertaken for the Department of Environment and Heritage by CSIRO Marine Research.](#)

Summary: This report is the final report of a two year study designed to identify and rank introduced marine species found within Australian waters (potential domestic target species) and those that are not found within Australian waters (potential international target species).

Available from: <http://www.marine.csiro.au/crimp/reports/PriorityPestsFinalreport.pdf> [Accessed 25 May 2005]

General information

Ahl, R. S., and S. P. Moss. 1999. Status of the nonindigenous crab, *Hemigrapsus sanguineus*, at Greenwich Point.... Northeastern Naturalist 6(3): 221-224.



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Hemigrapsus sanguineus*

[Bourdeau, P.E. and O Conner N.J. 1999. Prey preferences of the recently introduced Western Pacific Shore Crab, *Hemigrapsus sanguineus*, feeding on molluscs and macroalgae in Southeastern Massachusetts. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999, Massachusetts Institute of Technology, Cambridge, MA](#)

Summary: Prey of different sizes were offered to the newly introduced crab to see which they preferred.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs1.pdf> [Accessed Sept 20 2004]

[Brandhagen, M.D.; Ellis, D.J.; O Conner, N.J. and Tholleson, M. 1999. Geographic differentiation of an introduced crab species \(*Hemigrapsus sanguineus*\) on the Atlantic coast of North America. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999, Massachusetts Institute of Technology, Cambridge, MA](#)

Summary: An attempt to find out the origins of the invading crabs.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs1.pdf> [Accessed Sept 20 2004]

[Breton G., M. Faasse, P. Noel, and T. Vincent. 2002. A new alien crab in Europe. *Hemigrapsus sanguineus* \(Brachyurea: Grapsidae\). Journal of crustacean biology 22\(1\): 184-189.](#)

[Brousseau, D. J., A. Filipowicz, and J. A. Baglivo. 2001. Laboratory investigations of the effects of predator sex and size on prey selection by the Asian crab, *Hemigrapsus sanguineus*. Exp. Mar. Biol. Ecol.. 2001 Jul 30;262\(2\):199-210.](#)

[Brousseau, D. J., K. Kriksciun, and J. A. Baglivo. 2003. Fiddler crab burrow usage by teh Asian crab, *Hemigrapsus sanguineus*, in a Long Island Sound salt marsh. Northeastern Naturalist 10\(4\):415-420.](#)

[Brousseau, D.; Korchari, P. and Pflug, C. 1999. Food preference studies of the Japanese Shore Crab \(*Hemigrapsus sanguineus*\) from Western Long Island Sound. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999, Massachusetts Institute of Technology, Cambridge, MA](#)

Summary: An analysis of the diet preferences of the newly invasive shore crab.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs1.pdf> [Accessed Sept 20 2004]

[Casanova, T. 1999. The ecology of the Japanese Shore Crab \(*Hemigrapsus sanguineus* de Haan\) and its niche relationship to the Green Crab \(*Carcinus maenas* along the coast of Connecticut, U.S.A. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999, Massachusetts Institute of Technology, Cambridge, MA](#)

Summary: This newly arrived crab appears to be displacing another invasive crab.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs2.pdf> [Accessed Sept 20 2004]

[Casanova, T. 2001 The ecology of the Japanese Shore Crab \(*Hemigrapsus sanguineus* De Haan\) and its niche relationship to the Green Crab \(*Carcinus maenas* Linneus\) Along the coast of Connecticut, U.S.A. In Abstracts: Second International Conference on Marine Bioinvasions, March 9-11, 2001, New Orleans, LA](#)

Summary: The Japanese shore crab was found to be having a negative impact on the green crab population.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/2001/MBI2001abs2.pdf> [Accessed Sept 20 2004]

[DeGraaf, J.D. & Tyrrell, M.C.. \(2004\). Comparison of the feeding rates of two introduced crab species, *Carcinus maenas* and *Hemigrapsus sanguineus*, on the blue mussel, *Mytilus edulis*. 11. 163-166.](#)

[Epifanio, C.E., Park, S., Grey, E.K. and Waidner, L.A. 2003. Interactions between the asian shore crab \(*Hemigrapsus sanguineus*\) and the common mud crab \(*Panopeus herbstii*\): Larval supply vs post-settlement competition. In Abstracts: Third International Conference on Marine Bioinvasions, March 16-19, 2003, Scripps Institution of Oceanography La Jolla, California](#)

Summary: The displacement of the native common mud crab by the invasive asian shore crab maybe due to competition at a post-larval stage.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/2003/MBI2003abs3.pdf> [Accessed Sept 20 2004]

[Gerard, V. A., R. M. Cerrator, and A. A. Larson. 1999. Potential impacts of a western Pacific grapsid crab on intertidal communities of the northwestern Atlantic Ocean. Biological Invasions 1: 353-361, 1999.](#)

[Jensen, G. C., P. S. McDonald, D. A. Armstrong. 2002. East meets west: competitive interactions between green crab *Carcinus maenas*, and native and introduced shore crab *Hemigrapsus* spp. Mar Ecol Prog Ser Vol. 225: 251-262.](#)

[Kraemer, G.P. and Sellberg, M. 2001. Effects of the non-native crab *Hemigrapsus sanguineus* on crab community diversity at Edith Reed Sanctuary \(Rye, NY\) from 1998-2000. In Abstracts: Second International Conference on Marine Bioinvasions, March 9-11, 2001, New Orleans, LA](#)

Summary: Report into the expansion and increase in Japanese shore crab abundances.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/2001/MBI2001abs6.pdf> [Accessed Sept 20 2004]

[Larson, A.; Gerard, V. and Cerrato, R. 1999. Potential impact of the recently introduced Asian Shore Crab, *Hemigrapsus sanguineus*, on rocky intertidal communities of the Northeastern U.S. coast. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999, Massachusetts Institute of Technology, Cambridge, MA](#)

Summary: May have a large impact on another introduced species through predation.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs7.pdf> [Accessed Sept 20 2004]

[Ledesma, M. E., and N. J. Connor. 2001. Habitat and Diet of the Non-Native Crab *Hemigrapsus sanguineus* in Southeastern New England. Northeastern Naturalist: Vol. 8, No. 1, pp. 63-78.](#)

[Lohrer, A. M., and R. B. Whitlatch. 2002. Interactions among aliens: Apparent replacement of one exotic species by another. Ecology. 83\(3\). March, 2002. 719-732.](#)

[Lohrer, A.M.; Whitlatch, R.B.; Wada, K. and Fukui, Y. 1999. Patterns and predictability of biological invasions: A theoretical framework and case study using the Western Pacific Shore Crab *Hemigrapsus sanguineus*. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999, Massachusetts Institute of Technology, Cambridge, MA](#)

Summary: Development of a theoretical framework for the prediction of the effects of invasive species and the application of this framework on *H. sanguineus*.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs7.pdf> [Accessed Sept 20 2004]

[Mcdermott, J. J. 1998. The western Pacific brachyuran \(*H. sanguineus*: Grapsidae\), in its new habitat along the Atlantic coast of the United States: geographic distribution and ecology. ICES Journal of Marine Science, 55: 289-298. 1998](#)



GLOBAL INVASIVE SPECIES DATABASE

FULL ACCOUNT FOR: *Hemigrapsus sanguineus*

McDermott, J.J. 1999. A review of the natural history of the Asian Shore Crab *Hemigrapsus sanguineus* in the Western Atlantic with additional information on its biology. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999. Massachusetts Institute of Technology, Cambridge, MA

Summary: Study into the size and abundance of *H. sanguineus* on the Atlantic coast of America.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs8.pdf> [Accessed Sept 20 2004]

O Conner, N.J. 2001. The Asian Shore Crab *Hemigrapsus sanguineus* In New England: Changes in resident crab populations? In Abstracts: Second International Conference on Marine Bioinvasions, March 9-11, 2001. New Orleans, LA

Summary: The Asian shore crab and its influences on local and established populations of crabs.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/2001/MBI2001abs8.pdf> [Accessed Sept 20 2004]

O Conner, N.J., Bourdeau, P.E. and Ledesma, M.E. 1999. Northward extension of the geographic range of *Hemigrapsus sanguineus* in Massachusetts, 1996-1998. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999. Massachusetts Institute of Technology, Cambridge, MA

Summary: Paper documenting the spread of *H. sanguineus*.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs9.pdf> [Accessed Sept 20 2004]

Park, S. 1999. Early life history of *Hemigrapsus sanguineus*, a nonindigenous crab, in the Middle Atlantic Bight (USA). In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999. Massachusetts Institute of Technology, Cambridge, MA

Summary: Investigation into the juvenile stage of the *H. sanguineus*.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs9.pdf> [Accessed Sept 20 2004]

Percival, S.R. and Wilson, K. 2001. Feeding and substrate preferences of the Japanese Shore crab, *Hemigrapsus sanguineus*. In Abstracts: Second International Conference on Marine Bioinvasions, March 9-11, 2001. New Orleans, LA

Summary: Investigation into the substrate preferences and feeding behaviour of the invasive crab.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/2001/MBI2001abs8.pdf> [Accessed Sept 20 2004]

Richards, B. and Jensen, B. 1999. The incidence of *Hemigrapsus* relative to salinity values in the Delaware estuary and the inland bays of Delaware. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999. Massachusetts Institute of Technology, Cambridge, MA

Summary: Attempt to explain why *Hemigrapsus* has established in a large American estuary but not a smaller estuary.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs10.pdf> [Accessed Sept 20 2004]

Schubart, C. D. 2003. The East Asian shore crab *Hemigrapsus sanguineus* (Brachyure: Varunidae) in the Mediterranean Sea: an independent human-mediated introduction. *Sci. Mar.* 67(2) 195-200.

Tyrrell, M.C. 1999. Predicted impacts of the introduced crab, *Hemigrapsus sanguineus*, in New Hampshire. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999. Massachusetts Institute of Technology, Cambridge, MA

Summary: Attempt to model the invasion of *H. sanguineus* in New Hampshire.

Available from <http://massbay.mit.edu/resources/pdf/MarinePDF/1999/MarineAbs11.pdf> [Accessed Sept 20 2004]

Whitlatch, R.B. and Osman, R.W. 1999. Geographical distributions and organism-habitat associations of shallow-water introduced marine fauna in New England. In Abstracts: First National Conference on Marine Bioinvasions, January 24 -27, 1999. Massachusetts Institute of Technology, Cambridge, MA

Summary: Report into the distribution of some invasive benthic species in New England, USA

Available from: http://massbay.mit.edu/publications/marinebioinvasions/mbi1_abstracts.pdf [Accessed 8 February 2008]

World Register of Marine Species (WoRMS), 2011. *Hemigrapsus sanguineus* (De Haan, 1835)

Summary: Available from: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=158417> [Accessed 11 February 2011]