

FULL ACCOUNT FOR: Euglandina rosea



System: Terrestrial

Kingdom	Phylum	Class	Order	Family
Animalia	Mollusca	Gastropoda	Stylommatophora	Spiraxidae

Rosige Wolfsschnecke (German), rosy wolf snail (English), cannibal snail Common name

(English)

Synonym

Similar species

Summary The carnivorous rosy wolfsnail Euglandina rosea was introduced to Indian and

Pacific Ocean Islands from the 1950s onwards as a biological control agent for the giant African snail (Achatina fulica). E. rosea is not host specific meaning that native molluscs species are at risk of expatriation or even extinction if this mollusc-eating snail is introduced. Partulid tree snails of the French Polynesian Islands were particularly affected; having evolved separately from each other in isolated valleys, many Partulid tree snails have been lost and

today almost all the survivors exist only in zoos.

view this species on IUCN Red List

Species Description

The shell is large (up to 76 mm in height, 27.5 mm in diameter), thick and has prominent growth lines (University of Florida 2009). The shape of the shell is fusiform with a narrow ovate-lunate aperture and a truncated columella; typically, the shell color is brownish-pink (University of Florida 2009). Adult Euglandina grow from about seven to 10 cm long (Clifford et al. 2003).

Habitat Description

Euglandina rosea is usually found singly in hardwood forests, roadsides and urban gardens in its native range in Florida (Hubricht 1985, University of Florida 2009).

Reproduction

Euglandina rosea is a cross-fertilising egg-laying hermaphrodite. Chiu and Chou (1962, in Univeristy of Florida 2009) gave details of the biology of Euglandina in Taiwan. Individuals live up to 24 months. 25 to 35 eggs are laid in a shallow pocket in the soil. These hatch after 30 to 40 days.

Nutrition

Euglandina rosea feeds on other snails and slugs, which they track down by following the slime trails left by their prey (Clifford et al. 2003). It appears to prefer smaller individuals, which it swallows whole, but will attack large snails by entering through the shell aperture.



FULL ACCOUNT FOR: Euglandina rosea

General Impacts

Molluscs are the group most affected by extinction according to the 2007 International Union for Conservation of Nature (IUCN) Red List (Regnier 2009). The Pacific region has a wide diversity of mollusc species, most of them unique to the region, and the majority endemic to single islands or archipelagos (Cowie 1996 1997a, in Cowie and Cook 2001). More and more, these unique species are becoming replaced with a homogenous group of tropical tramp snail and slug species that are increasingly widespread (Cowie 1998a, R.H. Cowie, unpub., in Cowie and Cook 2001). Of the 400 extinct species we listed from oceanic islands, 234 lived on islands to which Euglandina rosea had been introduced, and it is highly probable that of these 234 extinctions, 134 (>50%) of them were ultimately caused by the introduction of *E. rosea* (Regnier et al. 2009).\r\n E. rosea contributed to the extinctions of endemic Partula tree snails in French Polynesia; the snails are widely distributed on most of the high islands of the tropical Pacific, except for the Hawaiian Islands (Murray et al. 1989, Cowie 1992, Hopper and Smith 1992, in Cowie and Cook 2001). E. rosea also contributed to the marked decline of endemic land snail fauna in Hawaii and Mauritius (Murry et al. 1988; Clarke et al. 1984; Hadfield 1986, Murray et al. 1988, Griffiths et al. 1993, Wells 1995, in Satoshi 2003). The best documented cases are those of the achatinelline tree snails, which are endemic to the Hawaiian Islands (Hadfield 1986, Hadfield et al. 1993). The native species mentioned seem especially vulnerable to heightened levels of predation because of their extremely slow rate of reproduction (Cowie 1992; Hadfield et al. 1993, in Cowie and Cook 2001).\r\n The carnivorous snail was introduced to control numbers of the giant African land snail (Achatina fulica) (Nishida and Napompeth 1975, in Cowie 2000). However, no rigorous scientific evidence exists that E. rosea controls A. fulica (Christensen 1984, in Cowie 2000) and, as a consequence, the World Conservation Union (IUCN) has formally condemned the deliberate introduction of E. rosea and other carnivorous snails. Most governments and other authorities appear to be aware of the potential threat posed to native fauna by E. rosea, however, under pressure to do something about A. fulica, they may misguidedly consider the introduction of E. rosea (and other species such as the flatworm *Platydemus manokwari*).

<u>Disease transmission</u>: *E. rosea* was found experimentally to be able to serve as both an intermediate and a paratenic host of *Angiostrongylus cantonensis*.



FULL ACCOUNT FOR: Euglandina rosea

Management Info

For a detailed account of the environmental impacts of *Euglandina rosea* please read: *Euglandina rosea* (Rosy Wolfsnail) Management Information. The information in this document is summarised below.

The future for some of French Polynesia's partulids may not be as bleak as once thought; according to recent studies relatively high genetic diversity is represented among living taxa and it may still be possible to preserve a representative sub-sampling of Raiatea and Tahiti's tree snail diversity (Lee *et al.* 2009; Ó Foighil 2009). Physical Control: The ultimate objective of captive breeding programs is the reintroduction of viable populations of endangered species into their natural habitats (Coote *et al.* 2004). Small exclosures have been built in Hawai'l and on Moorea (French Polynesia) to protect native tree snails from attack by *Euglandina rosea*. Legislation: It is almost impossible to prevent the within-island spread of *Euglandina* in French Polynesia (Coote *et al.* 1999). Between-island spread of *Euglandina* should be prevented by legislation. The Marqueses Islands, the Southern Cooks and the Australs provide refuges for some of the remaining partulid species (Lee *et al.* 2007a) and should be kept *Euglandina*-free. *E. rosea* is now legally considered to be a noxious species in French Polynesia; the introduction of live specimens and their transport from one island to another is forbidden (Meyer 1998).

Other: Since 1986 partulid snails have been the subject of an international breeding programme; the International Partula Conservation Programme manages a breeding programme for 25 species in 15 zoos worldwide. Introducing Society Island partulids to the Austral Islands that are free of the predator might ensure their long-term survival in the wild (Ó Foighil 2009). Coote & Loeve (2003) concluded that *E. rosea* was extinct in the wild on Huahine, strongly suggesting that the successful re-introduction of partulids into the wild on Huahine might be possible.

Conservation actions in the wild may be limited to identifying and protecting populations of partulid snails that offer some possibility of persistence in the presence of *Euglandina* (Ó Foighil 2009). Based on laboratory behavioral studies of the effect of temperature on *E. rosea* movement, Gerlach (1994, in Ó Foighil 2009) hypothesised that an altitudinal refuge above 600 to 700 m would exist for Society Island partulids. Research and Knowledge: Further research into the biology of *E. rosea*, and particularly its population dynamics, needs to be carried out. There are no known natural predators, so a species-specific toxin in snail bait, as tested in Hawaii (M. G. Hadfield pers. comm., in Coote *et al*, 1999), could be a promising approach. A good relationship between the Pacific Island Land Snail Group (PILSG) and the French Polynesian government authorities has developed, and joint initiatives for conservation and research are being planned (Coote *et al*. 1999). Education and Knowledge: Despite the lack of evidence supporting *Euglandina* as a successful biological control agent and despite the abundant evidence of their negative predatory impact on native snail fauna, carnivorous snail introductions continue (Cowrie 1992). Clearly public education about the French Polynesia's precious natural fauna and the dangers posed to such fauna by carnivorous biological control agents could help to reduce the likelihood of *Euglandina* being purposefully translocated to new islands. Local willingness and experience are already in place to conserve French Polynesia's partulid snails (Coote & Loeve 2003).

Principal source:

Compiler: IUCN SSC Invasive Species Specialist Group

Updates with support from the Overseas Territories Environmental Programme (OTEP) project XOT603, a joint project with the Cayman Islands Government - Department of Environment

Review: Review of updates under progress

Dr. Robert H. Cowie, Center for Conservation Research and Training, University of Hawaii, Honolulu, Hawaii

Pubblication date: 2010-02-17

ALIEN RANGE

[2] AMERICAN SAMOA [1] BAHAMAS

[1] BERMUDA [8] FRENCH POLYNESIA

[1] GUAM
Global Invasive Species Database (GISD) 2025. Species profile Euglandina rosea. Available from: https://iucngisd.org/gisd/species.php?sc=92 [Accessed 14 December 2025]



FULL ACCOUNT FOR: Euglandina rosea

[2] INDIA[3] JAPAN

[1] MADAGASCAR

[1] MAYOTTE

[1] NORTHERN MARIANA ISLANDS

[1] PAPUA NEW GUINEA

[1] SEYCHELLES [1] SRI LANKA

[1] UNITED STATES

[1] WALLIS AND FUTUNA

[1] INDONESIA

[2] KIRIBATI

[1] MAURITIUS

[1] NEW CALEDONIA

[1] PALAU

[1] REUNION

[1] SOLOMON ISLANDS

[1] TAIWAN

[1] VANUATU

Red List assessed species 72: EX = 43; EW = 10; CR = 12; EN = 1; VU = 2; DD = 4;

Achatinella mustelina CR
Partula affinis CR
Partula atilis EX
Partula auriculata EX
Partula callifera EX
Partula cedista EX
Partula clara CR
Partula clara CR
Partula cuneata EX
Partula dentifera EW
Partula dolorosa EX
Partula exigua EX
Partula filosa EX
Partula fusca EX
Partula hebe EW

Partula imperforata EX
Partula leptochila EX
Partula levistriata EX
Partula lutea EX
Partula mooreana EW
Partula otaheitana CR
Partula planilabrum EX
Partula protracta EX
Partula remota EX
Partula rosea EW
Partula sagitta EX
Partula taeniata CR
Partula tohiveana EW

Partula turgida EX
Partula varia EW
Samoana abbreviata CR
Samoana attenuata CR
Samoana conica EN
Samoana diaphana DD
Samoana hamadryas CR
Samoana jackieburchi EX

Erinna newcombi VU
Partula arguta EX
Partula aurantia EX
Partula bilineata EX
Partula candida EX
Partula citrina EX
Partula crassilabris EX
Partula cytherea EX
Partula dolichostoma EX
Partula eremita EX
Partula faba EW

Partula dolichostoma E
Partula eremita EX
Partula faba EW
Partula formosa EX
Partula garretti EX
Partula hyalina VU
Partula labrusca EX
Partula levilineata EX
Partula lugubris EX
Partula mirabilis EW
Partula navigatoria EX
Partula ovalis EX
Partula producta EX

Partula producta EX
Partula radiata EX
Partula robusta EX
Partula rustica EX
Partula suturalis EW
Partula thalia EX
Partula tristis EW
Partula umbilicata EX
Partula vittata EX
Samoana annectens DD
Samoana burchi CR
Samoana decussatula CR
Samoana ganymedes DD
Samoana inflata EX

Samoana magdalinae DD

Samoana oreas CR Samoana strigata CR BIBLIOGRAPHY

30 references found for Euglandina rosea

Managment information

Carlton, J.T. & G.M. Ruiz. 2003. Invasive species: vectors and management strategies. Island Press.



FULL ACCOUNT FOR: Euglandina rosea

Civeyrel, L. and Simberloff, D. 1996. A tale of two snails: is the cure worse than the disease? Biodiversity and Conservation 5: 1231-1252. **Summary:** A discussion of the introduction of predatory snails (notably *Euglandina rosea*), in putative attempts to control *A. fulica*. The devastating consequences on native land snail diversity, especially in the islands of the Pacific.

Clifford, K.T., L. Gross, K. Johnson, K.J. Martin, N. Shaheen & M.A. Harrington. 2003. Slime-Trail Tracking in the Predatory Snail, *Euglandina rosea, Behavioral Neuroscience* 117(5): 1086-1095.

Coote, T., D. Clarke, C.S. Hickman, J. Murray & P. Pearce-Kelly. 2004. Experimental Release of Endemic Partula Species, Extinct in the Wild, into a Protected Area of Natural Habitat on Moorea, *Pacific Science* 58(3): 429-434.

Cowie, R. H. 2000. Non-indigenous land and freshwater molluscs in the islands of the Pacific: conservation impacts and threats. In G. Sherley (ed.) Invasive species in the Pacific: a technical review and regional strategy. South Pacific Regional Environment Programme, Samoa: 143-172.

Summary: Discusses the conservation related impacts of the introduction of alien land and freshwater snails and slugs to the islands of the Pacific. Provides details of the main alien species of concern, identifies islands most at risk and islands on which to focus conservation efforts. Lists distribution details for all alien snails and slugs in the Pacific.

Gargominy, O. 2008. Beyond the alien invasion: a recently discovered radiation of Nesopupinae (Gastropods: Pulmonata: Vertiginidae) from the summits of Tahiti (Society Islands, French Polynesia), *Journal of Conchology 39*(5).

Hadfield, M. G., Miller, S. E. and Carwile, A. H. 1993. The decimation of endemic Hawai in tree snails by alien predators. American Zoologist 33: 610-622.

Summary: Discusses the impacts of alien rats and Euglandina rosea on native Hawaiian tree snails.

IUCN/SSC Invasive Species Specialist Group (ISSG)., 2010. A Compilation of Information Sources for Conservation Managers.

Summary: This compilation of information sources can be sorted on keywords for example: Baits & Lures, Non Target Species, Eradication, Monitoring, Risk Assessment, Weeds, Herbicides etc. This compilation is at present in Excel format, this will be web-enabled as a searchable database shortly. This version of the database has been developed by the IUCN SSC ISSG as part of an Overseas Territories Environmental Programme funded project XOT603 in partnership with the Cayman Islands Government - Department of Environment. The compilation is a work under progress, the ISSG will manage, maintain and enhance the database with current and newly published information, reports, journal articles etc.

Mead, A. R. 1961. The giant African snail: a problem in economic malacology. Chicago, University of Chicago Press.

Summary: Major treatise on the worldwide spread of A. fulica, its impacts, and management.

Murray, J., Murray, E., Johnson, M. S. and Clarke, B. 1988. The extinction of Partula on Moorea. Pacific Science 42: 150-153.

Summary: Reports the final demise of all seven *Partula* species of Moorea in the face of the spread of *E. rosea* and the imminent threat to *Partula* on Tahiti.

Varnham, K. 2006. Non-native species in UK Overseas Territories: a review. JNCC Report 372. Peterborough: United Kingdom.

Summary: This database compiles information on alien species from British Overseas Territories.

Available from: http://www.jncc.gov.uk/page-3660 [Accessed 10 November 2009]

General information

Campbell, B.G. and Little, M.D. 1988. The finding of *Angiostrongylus cantonensis* in rats in New Orleans, *Am J Trop Med Hyg. 38*(3): Coote, T. and Loeve, E. 2003. From 61 species to five: endemic tree snails of the Society Islands fall prey to an ill-judged biological control programme, *Oryx 37*(1): abstract.

Coote, Trevor, Zoological Society of London. Field Report 2000/2001 unpub.

Cowie, R.H. and Cook, R.P. 2001. Extinction or survival: partulid tree snails in American Samoa, *Biodiversity and Conservation 10*(2). Gargominy, O., Bouchet, P., Pascal, M., Jaffre, T. and Tourneu, J. C. 1996. Consequences des introductions d'especes animals et vegetales sur la biodiversite en Nouvelle-Caledonie. Rev. Ecol. (Terre Vie) 51: 375-401.

Summary: Consequences to the biodiversity of New Caledonia of the introduction of plant and animal species.

Gargominy, O. (Ed.). 2003. Biodiversit vet conservation dans les collectivit vs fran aises d outre-mer. Comit fran ais pour l'UICN, Paris.

Summary: Synth se sur la biodiversit des se les frantaises d'outre-mer et les enjeux de conservation.

Available from: http://www.uicn.fr/Biodiversite-outre-mer-2003.html [Accessed 26 March 2008]

Griffiths, O., Cook, A. and Wells, S. M. 1993. The diet of the carnivorous snail *Euglandina rosea* in Mauritius and its implications for threatened island gastropod faunas. Journal of Zoology 229: 79-89.

Summary: Assesses the feeding prefernces of *E. rosea*, showing that it preferes snails other than *A. fulica*, and preferes small over large snails. Lists the regions into which it has been introduced and the impacts in those regions.

Griffiths, O. Cook, A. and Wells, S.M. 1993. The diet of the introduced carnivorous snail *Euglandina rosea* in Mauritius and its implications for threatened island gastropod faunas, *Journal of Zoology* 229(1): abstract.

Griffiths, O. Florens, V. 2006. A field guide to the non-marine molluscs of the Mascarene Islands (Mauritius, Rodrigues and Réunion) and the Northern Dependencies of Mauritius. Bioculture Press, Mauritius

Hopper, D.R. and Smith, B.D. 1992. Status of tree snails (Gastropoda: Partulidae) on Guam, with a resurvey of sites studied by H.E. Crampton in 1920, *Pacific Science 46*(1): abstract.

Howarth, F.G. 1991. Environmental Impacts of Classical Biological Control, Annu. Rev. Entomol. 36.

ITIS (Integrated Taxonomic Information System), 2005. Online Database Euglandina rosea

Summary: An online database that provides taxonomic information, common names, synonyms and geographical jurisdiction of a species. In addition links are provided to retrieve biological records and collection information from the Global Biodiversity Information Facility (GBIF) Data Portal and bioscience articles from BioOne journals.

Available from:

http://www.cbif.gc.ca/pls/itisca/taxastep?king=every&p_action=containing&taxa=Euglandina+rosea&p_format=&p_ifx=plglt&p_lang=[Accessed March 2005]

Kinzie, R.A. 1992. Predation by the introduced carnivorous snail *Euglandina rosea* (Ferussac) on endemic aquatic lymnaeid snails in Hawaii, *Biological Conservation 60*(3): abstract.



FULL ACCOUNT FOR: Euglandina rosea

Lee, T., J-Y. Meyer, J.B. Burch, P. Pearce-Kelly & D. O Foighil. 2008. Not completely lost: two partulid tree snail species persist on the highest peak of Raiatea, French Polynesia, Fauna & Flora International, Oryx, 42(4): 615-619.

Mus vum national d Histoire naturelle [Ed]. 2003-2006. Euglandina rosea (Forussac, 1821) Inventaire national du Patrimoine naturel Summary: Base de données en ligne sur le patrimoine naturel franéais.

Available from:

http://inpn.mnhn.fr/isb/servlet/ISBServlet?action=Espece&typeAction=10&pageReturn=ficheEspeceDescription.jsp&numero_taxon=433018 [Accessed 1 April 2008]

Ohbayashi, T., I. Okochi, H. Sato, T. Ono & S. Chiba. 2007. Rapid decline of endemic snails in the Ogasawara Islands, Western Pacific Ocean, *Appl. Entomol. Zool.* 42(3): 479-485.

Regnier, C., B. Fontaine & P. Bouchet. 2009. Not Knowing, Not Recording, Not Listing: Numerous Unnoticed Mollusk Extinctions, Conservation Biology 23(5): 1214 - 1221.

Satoshi, C. 2003. Species Diversity and Conservation of Mandarina, an Endemic Land Snail of the Ogasawara Islands.

Summary: Available from: http://www.airies.or.jp/publication/ger/pdf/07-01-04.pdf [Accessed 4 March 2006] University of Florida. 2009. Snails Eating Snails of Florida (Euglandina rosea (Ferussac 1821) (Gastropoda: Spiraxidae), Rumina decollata (Linnaeus 1758) (Gastropoda: Subulinidae), Haplotrema concavum (Say 1821) (Gastropoda: Haplotrematidae), Gulella bicolor (Hutton 1834)

(Gastropoda: Streptaxidae), Varicella gracillima floridana Pilsbry 1907)

Summary: Available from: http://edis.ifas.ufl.edu/in523 [Accessed 24 February 2010]