

MV (Massive) *Pelophylax bergeri*

| | |
|---|--|
| Date assessed | 2021-10-18 |
| Year published | 2021 |
| Eicat category | MV (Massive) |
| Justification for EICAT assessment | Impact through hybridisation between <i>P. bergeri</i> and <i>P. lessonae</i> is massive, hybrids have completely replaced the native <i>P. lessonae</i> (Dubey et al. 2014, Dufresnes et al. 2007, Dufresnes and Dubey 2020). The deep introgression has collapsed the genetic structure of the native <i>Pelophylax</i> populations (Dubey et al. 2014). |
| Confidence rating | High |
| Mechanism(s) of maximum impact | Hybridisation |
| Countries of most severe impact | Switzerland; France |
| Description of impact | Hybridisation - genetic introgression between <i>Pelophylax bergeri</i> and the native species <i>P. lessonae</i> and <i>P. esculentus</i> has led to the replacement of most pure native populations. |
| Assessor | F. André de Villiers |
| Contributors | Carla Wagener; Mohlamatsane Mokhatla; James Baxter-Gilbert; Corey Thorp; Alexander D. Rebelo; Sarah J. Davies; Nitya Prakash Mohanty; John Measey; Giovanni Vimercati; Sabrina Kumschick; Khensani Nkuna |
| Reviewers | EICAT authority |
| Recommended citation | F. André de Villiers. (2026). <i>Pelophylax bergeri</i> . IUCN Environmental Impact Classification for Alien Taxa (EICAT) . |

